

Amblyomma variegatum

Tropical Bont Tick, Tropical
African Bont Tick

Last Updated: September 2009



IOWA STATE UNIVERSITY
College of Veterinary Medicine



- OIE Collaborating Centres:
- Diagnosis of Animal Disease and Vaccine Evaluation in the Americas
 - Day-One Veterinary Competencies and Continuing Education



www.cfsph.iastate.edu
Email: cfsph@iastate.edu

Importance

Amblyomma variegatum is a hard tick that feeds on a number of domesticated animals including cattle, sheep, goats, horses and dogs, as well as humans. The long mouthparts of *A. variegatum* leave large wounds, and make this tick difficult to remove manually. Its bite is severe and painful, and can result in significant damage to the skin. Secondary infections can cause septic wounds or abscesses, and inflammation on the teats of cows may affect milk production. In some regions, *Amblyomma* bite wounds may become infested by screwworms. In addition, *A. variegatum* is a host for a number of microbial pathogens including *Ehrlichia ruminantium* (formerly *Cowdria ruminantium*), the agent of heartwater, and *Rickettsia africae*, the agent of African tick-bite fever, which is an emerging zoonosis in rural sub-Saharan Africa and the Caribbean.

Species Affected

Immature ticks feed on small mammals, ground-feeding birds, reptiles, cattle, sheep and goats. Adult ticks prefer cattle, but can also be found on sheep, goats, horses, camels, dogs and some large wildlife including antelope.

Geographic Distribution

A. variegatum is found in the tropics and subtropics. This tick is endemic in savannas in many countries in sub-Saharan Africa, as well as in southern Arabia, the Caribbean, and some islands in the Atlantic and Indian Oceans. An eradication program is in progress in the Caribbean; St. Kitts, St. Lucia, Montserrat, Anguilla, Barbados and Dominica qualified for 'provisionally free' certification by 2002, although St. Kitts was re-infested in 2004.

Life Cycle

A. variegatum is a three-host tick. Immature ticks feed on small mammals, ground-feeding birds and reptiles, as well as cattle, sheep and goats. Adult ticks prefer cattle, but can also be found on other livestock including camels, as well as dogs and some wildlife. The adult ticks are usually found on the relatively hairless parts of the body; most are located on the ventral body surface and the genitalia, or under the tail. Adult *A. variegatum* feed mainly in the rainy season, while the immature ticks feed primarily during the dry season.

Identification

 [Click to view images](#)

A. variegatum is a member of the family Ixodidae (hard ticks). Hard ticks have a dorsal shield (scutum) and their mouthparts (capitulum) protrude forward when they are seen from above. *Amblyomma* ticks are large ticks with long, strong mouthparts. The palps are long; the second segment is twice as long as it is wide. Eyes are present and the festoons are well developed. The males have no adanal shields, accessory shields or subanal shield.

Female *A. variegatum* are brown, but the males are brightly ornamented with orange. When they are engorged, the adult female ticks are about the size of a nutmeg. Tick identification to the species level can be difficult, and ticks should be submitted to an expert for identification whenever possible.

Recommended actions if *Amblyomma variegatum* is suspected

Notification of authorities

Known or suspected *A. variegatum* infestations should be reported immediately to state or federal authorities.

Federal: Area Veterinarians in Charge (AVIC):

https://www.aphis.usda.gov/aphis/ourfocus/animalhealth/contact-us/CT_Vet_Acred_Asst_DD

State Animal Health Officials:

http://www.usaha.org/upload/Federal%20and%20State%20Health/StateAnimalHealthOfficials_rev.pdf

Amblyomma variegatum

Control

Measures used to exclude exotic ticks from a country include pre-export inspection and certification that the animals are free of ectoparasites, quarantines upon entry, and treatment with acaricides. In countries where *A. variegatum* is already present, acaricides can eliminate the ticks from the animal, but do not prevent reinfestation and must be repeated periodically. Three-host ticks spend at least 90% of their life cycle in the environment rather than on the host animal; ticks in the environment must also be controlled to prevent their spread.

If ticks are already widespread in a region, eradication can be difficult. The Caribbean eradication program is based on animal identification and mandatory periodic acaricide treatment of livestock by farmers, as well as public education, surveillance, quarantines and movement restrictions.

Public Health

Amblyomma tick bites are painful, and the wound may become infected. In addition, *A. variegatum* can transmit exotic diseases including African tick bite fever to humans.

Internet Resources

Acarology WWW Home Page

http://www.nhm.ac.uk/hosted_sites/acarology/

Caribbean *Amblyomma* Programme

<http://www.fao.org/english/newsroom/audio/cap.pdf>

Food and Agriculture Organization of the United Nations (FAO). Ticks and Tick-borne Diseases.

<http://www.fao.org/docrep/004/X6538E/X6538E00.HTM>

The Merck Veterinary Manual

<http://www.merckvetmanual.com>

The University of Edinburgh. The Tick Collection.

<http://www.nhc.ed.ac.uk/index.php?page=24.25.121>

United States Animal Health Association. Foreign Animal Diseases.

http://www.aphis.usda.gov/emergency_response/downloads/nahems/fad.pdf

World Organization for Animal Health (OIE)

<http://www.oie.int>

OIE International Animal Health Code

<http://www.oie.int/international-standard-setting/terrestrial-code/access-online/>

Acknowledgements

This factsheet was written by Anna Rovid Spickler, DVM, PhD, Veterinary Specialist from the Center for Food Security and Public Health. The U.S. Department of Agriculture Animal and Plant Health Inspection Service

(USDA APHIS) provided funding for this factsheet through a series of cooperative agreements related to the development of resources for initial accreditation training.

The following format can be used to cite this factsheet.

Spickler, Anna Rovid. 2009. *Amblyomma variegatum*.

Retrieved from <http://www.cfsph.iastate.edu/DiseaseInfo/factsheets.php>.

References

- Food and Agriculture Organization of the United Nations [FAO]. Caribbean *Amblyomma* Programme (CAP) [online]. Available at: <http://www.fao.org/AG/AGAINFO/projects/en/cap/home.html>. * Accessed 11 Dec 2006.
- Kahn CM, Line S, editors. The Merck veterinary manual [online]. Whitehouse Station, NJ: Merck and Co; 2003. *Amblyomma* spp. Available at: <http://www.merckvetmanual.com/mvm/index.jsp?cfile=htm/bc/72107.htm>. Accessed 11 Dec 2006.
- Kelly PJ, Fournier PE, Parola P, Raoult D. A survey for spotted fever group rickettsiae and ehrlichiae in *Amblyomma variegatum* from St. Kitts and Nevis. *Am J Trop Med Hyg*. 2003;69:58-9.
- New South Wales Department of Agriculture. Identification of the paralysis tick *I. holocyclus* and related ticks [online]. New South Wales Department of Agriculture; 2001 Feb. Available at: <http://members.ozemail.com.au/~norbertf/identification.htm>. * Accessed 29 November 2001.
- Parola P, Inokuma H, Camicas JL, Brouqui P, Raoult D. Detection and identification of spotted fever group Rickettsiae and Ehrlichiae in African ticks. *Emerg Infect Dis*. 2001;7:1014-7.
- Pegram RG, Eddy C. Progress towards the eradication of *Amblyomma variegatum* from the Caribbean. *Exp Appl Acarol*. 2002;28:273-81.
- Pegram RG, Rota A, Onkelinx R, Wilson DD, Bartlette P, Nisbett BS, Swanston G, Vanterpool P, de Castro JJ. Eradicating the tropical bont tick from the Caribbean [online]. Available at: <http://www.fao.org/DOCREP/W2650T/w2650t06.htm>. Accessed 11 Dec. 2006.
- United States Animal Health Association [USAHA] Committee on Parasitic Diseases. Tropical bont tick eradication programs in the Caribbean [online]. In: USAHA 2005 Proceedings; 2005 Nov. 3-9; Hershey, PA. Available at: <http://www.usaha.org/committees/resolutions/2005/resolution49-2005.pdf>. * Accessed 11 Dec. 2006.
- Wilson DD, Bram RA. Foreign pests and vectors of arthropod-borne diseases. In: *Foreign Animal Diseases*. Richmond, VA: United States Animal Health Association; 1998. Available at: http://www.aphis.usda.gov/emergency_response/downloads/nahems/fad.pdf. Accessed 11 Dec 2006.

* Link is defunct