

# African Swine Fever

*Peste Porcine Africaine,  
Fiebre Porcina Africana,  
Pestis Africana Suum,  
Maladie de Montgomery,  
Warthog Disease,  
Afrikaanse Varkpes,  
Afrikanische Schweinepest*

**Last Updated:** June 2025



**IOWA STATE UNIVERSITY**  
College of Veterinary Medicine



## Importance

African swine fever (ASF) is an important viral disease of pigs that has spread extensively throughout the world since 2007 and has become a serious threat to pork production. ASF viruses range from highly pathogenic strains that may kill nearly the entire herd to less virulent isolates that cause a milder, nonspecific illness difficult to recognize as African swine fever. There is no effective treatment, and severely affected pigs usually die. The spread of ASF virus (ASFV) is facilitated by a number of factors, including its persistence for long periods in uncooked pork products, which may be fed to pigs in food scraps (pig swill), and its ability to become established in wild or feral suids. In some areas, control is complicated by the establishment of the virus in *Ornithodoros* ticks, which can act as biological vectors. One tick vector hindered eradication efforts during a previous outbreak in Spain and Portugal, where complete elimination of the virus took more than 30 years.

For many years, African swine fever was mostly confined to sub-Saharan Africa, where viruses from wild warthogs, thought to be the original host, had become widely established in domesticated pigs. In 2007, ASFV was accidentally introduced into the Caucasus region of Eurasia, most likely in pig swill. This highly virulent virus caused outbreaks on pig farms, became established in wild boar, and has since spread across much of Europe and Asia. It was recently introduced to the island of Hispaniola (Haiti and the Dominican Republic) in the Caribbean, and there is a possibility it might occur among wild boar in parts of the Middle East. The virus has also diversified, generating isolates of widely varying virulence, which are less likely to be recognized quickly. Controlling this outbreak has been difficult, with reports of recurrence in wild boar populations where ASFV was thought to have been eliminated, and a number of difficulties in developing an effective vaccine that does not revert to virulence.

## Etiology

African swine fever results from infection by African swine fever virus, an enveloped virus in the genus *Asfivirus* and family *Asfarviridae*. ASFV isolates differ greatly in virulence, ranging from highly pathogenic viruses that kill most pigs to strains that result only in seroconversion. At least 24 genotypes of ASFV have been identified, many from wildlife cycles in Africa. A number of these genotypes have also been found at some time in domesticated pigs, though only some are adapted to efficient transmission in these animals.

One genotype I virus has been endemic in Sardinia (Italy) since the 1960s, while the virus introduced into Eurasia in 2007 belongs to genotype II. Although the latter virus was highly virulent when introduced, its descendants have diversified, with isolates of varying virulence now circulating in Europe and Asia. In addition, a genotype I virus of low virulence, possibly introduced via an unauthorized vaccine, was identified in some herds in China and has recombined with genotype II viruses. At least some of these recombinants, which have since been found in other parts of Eurasia, are highly virulent.

## Species Affected

African swine fever affects members of the pig family (Suidae). Species known to be susceptible to infection include domesticated pigs and wild boar (both subspecies of *Sus scrofa*), warthogs (*Phacochoerus* spp.), bush pigs (*Potamochoerus larvatus* and *Potamochoerus porcus*), giant forest hogs (*Hylochoerus* spp.) and some wild Asian suids such as *Sus barbatus* and *Sus cebifrons*. Most of these animals can develop clinical signs, although infections in warthogs mostly seem to be subclinical or mild. Some older reviews and textbooks suggest that peccaries (*Tayassu* spp.) may become infected without clinical signs, although one attempt to infect collared peccaries (*Tayassu tajacu*) in 1969 was unsuccessful. Recent reviews state that peccaries are not susceptible to ASFV. Known maintenance hosts for the virus include domesticated pigs, wild boar and warthogs, while the role of other wild suids is unclear.

## Zoonotic potential

There is no evidence that ASFV infects humans.

## Geographic Distribution

African swine fever is endemic in much of sub-Saharan Africa and some nearby islands, e.g., Madagascar. Outbreaks were seen occasionally outside Africa before 2007, affecting parts of Europe, the Caribbean (Cuba, Hispaniola) and South America (Brazil). These viruses were all eventually eradicated, with the exception of a genotype I virus that has persisted for decades on the Mediterranean island of Sardinia (Italy).

A genotype II virus introduced into the Republic of Georgia in 2007 has since spread to domesticated swine and/or wild boars in many countries in Europe and Asia. In 2021, viruses from this outbreak became established in Hispaniola (Dominican Republic and Haiti), which is part of the Greater Antilles group of islands in the Caribbean. They were also found in wild boar in Iran in 2010, though there have been no reports of ASFV in the Middle East since then. Genotype I viruses, possibly introduced via the use of live attenuated vaccines, have been detected in some herds in China, and genotype I/II recombinants have since been found in China and other locations, including Vietnam and Russia.

Although some countries in the affected regions maintain an ASF-free domestic pig population, most reports of apparent virus elimination from wild boar in Europe have been followed by either recrudescence after low levels of virus circulation among boar (e.g., Estonia) or apparent reintroduction from neighboring countries (e.g., Czech Republic).

## Transmission

African swine fever can be transmitted either directly between animals or by the soft tick genus *Ornithodoros*, which acts as a biological vector. Domesticated pigs and wild boar can shed ASFV in most or all secretions and excretions including oronasal fluid, urine, feces and semen. While some pigs may begin to shed this virus 2 days before the onset of clinical signs, some studies suggest that the majority of virus shedding occurs late in the illness. Blood and tissues contain large amounts of virus, and massive environmental contamination may result if blood is shed during necropsies or pig fights, or if a pig develops bloody diarrhea. Although animals that recover can apparently clear the virus completely, some pigs may become chronically infected and continue to shed the virus. There is currently no evidence that viable ASFV persists long-term in a latent state in asymptomatic, healthy animals that have no evidence of viremia. Information about virus shedding in other suids is more limited; however, virus replication is much lower in adult warthogs, which do not seem to readily transmit the virus by direct contact.

Most pigs are thought to become infected with ASFV via inhalation or ingestion. Because this virus can persist in tissues after death, it can be spread by feeding uncooked or undercooked pig swill that contains tissues from infected animals. Cannibalism of dead animals might be important in some outbreaks. Fairly high doses of the virus are generally required to infect a pig in solid feed, though ingestion of the

virus in liquids seems to be more efficient. Inhalation requires only low viral doses. Aerosolized viruses may contribute to transmission within a building or farm, but current evidence suggests that this only occurs over relatively short distances. Boars can infect sows by artificial insemination as well as during natural breeding. Transplacental transmission has also been demonstrated; however, all of the infected fetuses reported to date either died *in utero* or shortly after birth.

*Ornithodoros* spp. ticks, which can infect suids either through bites or when they are eaten by the pig, are biological vectors for ASFV. In some parts of Africa, these viruses cycle between juvenile warthogs and soft ticks of the *Ornithodoros moubata* complex, which live in their burrows. Transstadial, transovarial and sexual transmission has been demonstrated in these ticks. A similar cycle can occur between domesticated pigs and the *Ornithodoros* spp. ticks that colonize their pens, and *O. erraticus* helped maintain ASFV on the Iberian Peninsula during the European outbreaks in the 1950s and 1960s. *Ornithodoros* spp. ticks are long-lived, and ASFV has been reported to persist in their colonies for several years (e.g., 5 years in *O. erraticus*). However, they can eventually clear the virus if they are not reinfected. Other blood-sucking insects such as various biting flies (e.g., *Stomoxys* spp., tabanids), mosquitoes, hard ticks or swine lice (e.g., *Haematopinus suis*) might be able to transmit ASFV mechanically for a short time, though their significance in transmission, if any, is still unclear. In addition, some potential mechanical vectors, such as swine lice, have limited mobility.

ASFV can also be spread on fomites, including vehicles, feed and equipment. Except in animal tissues, most studies of its persistence under controlled laboratory conditions, using various fomites including soil and unprocessed grains, have reported survival times ranging from less than a day to approximately 1-2 weeks, at temperatures from 4°C (39°F) to around 20°C (68°F). Survival was generally longer under cold conditions and when the virus was protected in feces or other organic matter. There are, however, a few outliers among these studies. One reported that ASFV was infectious for more than 16 weeks in sterilized soil samples at 4°C, and for either 2 or 3 weeks at 22°C (72°F), depending on whether or not sheep erythrocytes were added to the soil. Another study found that, in three different sterilized feed components, the virus remained viable for 5 days to 4 months at 4°C, 1 day to at least 21 days at 20°C, and < 1 day to at least 7 days at 35°C (95°F). Particularly long survival was found in sterilized soybean meal. Whether studies of virus persistence in sterilized materials accurately reflect real world conditions is uncertain, as the presence of other microorganisms can be detrimental to the survival of some viruses.

Studies on the persistence of ASFV in water, which might protect an enveloped virus from desiccation, reported that it remained viable in sterilized or distilled water for as long as 3-7 weeks at 22-24°C (72-75°F), and for at least 2 months either at 4-6°C (39-46°F) or when frozen at -16 to -20°C (-4 to 3°F). A study that used samples of river water found that ASFV remained viable for at least 42 days at 4°C,

28 days at 15°C (59°F), and 14 days at 21°C, but survived longer in distilled water. While these studies are likely to be applicable to water sources such as drinking troughs, their relevance to natural water sources such as rivers or lakes, where added viruses would be diluted to much lower levels, is unclear. The survival of ASFV in water appears to be reduced by the presence of ciliates but prolonged by freshwater snails. It has been reported to accumulate in the tissues of snails and leeches; however, feeding these snails to pigs did not result in infections, and the virus appeared to enter an abortive replication cycle in the leeches.

ASFV can persist for relatively long periods in refrigerated, frozen or preserved pig tissues, including a year and a half in blood and approximately 5 months in boned meat, both stored at 4°C, and 140 days in salted dried hams, with one study suggesting that it might persist for years in frozen carcasses. A study that used pork products made from experimentally infected animals detected live viruses in dry cured salami at 18 days but not 26 days after processing, in dry cured pork belly at 60 but not 137 days, and in dry cured loin at 83 but not 137 days. Studies that examined various porcine tissue and organ samples, generally as a proxy for virus survival in carcasses in the environment, reported that, depending on the substrate, ASFV persisted for periods ranging from a month to 2 years when frozen at -20°C or stored at 4°C, though infectious virus usually could not be found in tissues after 1-2 weeks at 20°C.

There is limited field data on ASFV survival outside the laboratory, where it might be destroyed more rapidly on fomites by sunlight and desiccation, and by rigor mortis and various decomposition processes in carcasses. Some studies found that pigs did not become infected when placed in pens that had housed animals with clinical signs 3-5 days earlier, if there was either no gross contamination with blood or any visible blood had been washed away and bloodstained areas decontaminated. In another report, researchers could not recover any infectious virus from pens 7 days after the removal of experimentally infected pigs with clinical signs that included bloody diarrhea. No infectious virus was found by three groups of researchers who examined the soil where naturally infected wild boar carcasses had been removed, looked for the virus in the bones of buried and unburied boar carcasses, or tested soil and organ samples from wild boar carcasses buried 18 days to 15 months earlier, though PCR tests detected viral nucleic acids in each of the 3 experiments. Viable ASFV was also reported to be absent from composted pig carcasses after 3 days.

### Disinfection

ASFV has been shown to be susceptible to a number of disinfectants including sodium hypochlorite, sodium hydroxide (caustic soda) and glutaraldehyde, as well as certain phenols, iodophors, peroxygens (e.g., potassium peroxymonosulfate), acids (citric, acetic), quaternary ammonium compounds and multi-component disinfectants such as Virkon™. As with other viruses, some disinfectants are more effective than others in the presence of significant

soiling, low environmental temperatures can affect the efficacy of certain agents, and prolonged contact time may be needed in some circumstances.

Heat is also effective in destroying this virus. Although 30 minutes at 60°C (140°F) is reported to be sufficient to inactivate ASFV in serum and body fluids, unprocessed meat must be heated to at least 70°C (158°F) for 30 minutes. Recommendations from different agencies for the heat inactivation of pig swill have varied. A recent study's theoretical calculations, based on observations of inactivation in small amounts of several different swills, suggest that inactivation may occur within a few minutes at 90°C (194 °F) but requires at least 2 hours at 70°C. Whether all of the swill will reach a given temperature must, however, also be considered. ASFV in serum-free medium can be inactivated by pH less than 3.9 or greater than 11.5.

### Incubation Period

The incubation period in domesticated pigs is reported to be 4-19 days in naturally-acquired cases and 2-21 days after experimental inoculation, with highly virulent viruses often affecting animals within a week. Some comparative studies suggest that wild boar might develop clinical signs more quickly than pigs after oronasal exposure.

### Clinical Signs

African swine fever can present as a peracute, acute, subacute or chronic illness. Some infected animals may also seroconvert without becoming ill. The course of the disease is generally correlated with the virulence of the virus; however, a given virus can cause more than one form, and some animals in a herd may remain asymptomatic or develop only transient signs while others become severely ill. Even in herds infected with highly virulent isolates, severely ill pigs are sometimes uncommon until the later stages of an outbreak, with most affected animals initially having mild, nonspecific clinical signs.

Pigs with peracute African swine fever die rapidly, usually with few or no clinical signs, while animals with acute cases typically have a high fever with nonspecific signs such as anorexia, lethargy, weakness and recumbency. Erythema may be evident, particularly in white animals, and some animals develop cyanotic skin blotching. The latter sign tends to be noted most often on the ears, snout, tail and perianal region, abdomen, lower legs and/or hams. Gastrointestinal signs including diarrhea, constipation, vomiting and signs of abdominal pain are also common. The diarrhea is initially mucoid and may later become bloody. Some pigs can also develop other hemorrhagic signs, such as epistaxis or cutaneous petechiae and ecchymoses. Respiratory signs (including dyspnea), nasal and conjunctival discharges are common in some outbreaks, and neurological signs have been reported. Pregnant animals frequently abort. Leukopenia and thrombocytopenia of varying severity may be detected in laboratory tests. Death often occurs within 7-10 days.

Subacute African swine fever is similar, but with less severe clinical signs. In some herds, abortions may be the first sign of these outbreaks. Thrombocytopenia and leukopenia may be transient in this form; however, petechiae or other hemorrhagic signs can sometimes be seen. Pigs with the subacute form of African swine fever usually die or recover within 3-4 weeks. In one outbreak, most sows that had recovered clinically but were still PCR positive either aborted or gave birth to stillborn and/or mummified fetuses. Some of these sows also produced live neonates with congenital tremors and ataxia that died within the first few days. Viral antigens and nucleic acids were found in the brains of the piglets with neurological signs, suggesting that African swine fever was the cause.

Chronically infected pigs usually have nonspecific signs such as an intermittent low fever, appetite loss and depression, and may be thin, emaciated and/or stunted. Some animals also cough or have other signs of respiratory issues, develop swollen joints, or have gastrointestinal signs such as diarrhea and/or occasional vomiting. Ulcers and reddened or raised necrotic skin foci may appear over body protrusions and other areas subject to trauma. Some of the clinical signs seen in these animals may be due to secondary infections. Chronically infected animals may eventually die.

Signs in experimentally infected wild boar are similar to those in domesticated pigs; however, some studies reported that they became ill more rapidly and died sooner. Skin lesions may also be absent or less evident in wild boar, with erythema usually limited to areas of thinner or hairless skin such as the abdomen and inguinal region. Some runted animals inoculated with very low viral doses had few or no clinical signs, including fever, before death. Warthogs and bush pigs usually become infected asymptotically or have mild cases.

### Post Mortem Lesions [Click to view images](#)

The gross lesions of African swine fever are highly variable, and are influenced by the virulence of the isolate and the course of the disease. Animals that die peracutely may have few or poorly developed lesions, while subacutely infected animals sometimes have more widespread organ involvement than those with acute cases.

The carcass is often in good condition in animals that die relatively soon after they become infected, but there may be bluish-purple discoloration and/or hemorrhages in the skin, and some pigs have external indications of bloody diarrhea or other internal hemorrhages. The major internal lesions are hemorrhagic, and can be seen most consistently in the spleen, lymph nodes, kidneys and heart. The spleen can be very large, friable, and dark red to black in some pigs; in others, it is enlarged but not friable, with a color closer to normal. The lymph nodes are often swollen and hemorrhagic, and may look like blood clots. Many reports indicate that the abdominal lymph nodes, including the gastrohepatic, mesenteric, ileocecal and renal nodes, are affected most often; however, the submandibular and medial retropharyngeal nodes also had similar lesions in some

animals exposed to the virus via the oronasal route. Petechiae are frequently noted on the cortical and cut surfaces of the kidneys, and sometimes in the renal pelvis, and perirenal edema may be evident. Hemorrhages, petechiae and/or ecchymoses may also be found in other organs including the urinary bladder, lungs, stomach and intestines, and pulmonary edema and congestion can be prominent in some pigs. Other reported lesions include congestion of the liver, edema in the wall of the gall bladder and bile duct, and straw-colored or blood-stained fluid in the pleural, pericardial and/or peritoneal cavities. In some cases, the brain and meninges are congested, edematous or hemorrhagic.

Lesions that have been noted in aborted fetuses in some reports include generalized edema (anasarca), a mottled liver, hemorrhagic tonsils, excess abdominal fluid, and petechiae or ecchymoses in the skin, myocardium and other organs. Petechiae may also be detected in the placenta. Five newborn piglets with congenital tremors had enlarged and hemorrhagic mandibular and inguinal lymph nodes, petechiae in the kidney, and congested meninges. One piglet also had hydranencephaly.

The carcasses of pigs with chronic African swine fever are often in poor condition, thin or emaciated. These animals may have a variety of lesions including focal areas of skin necrosis, skin ulcers, consolidated lobules in the lung, caseous pneumonia, nonseptic fibrinous pericarditis, pleural adhesions, generalized lymphadenopathy and swollen joints.

### Diagnostic Tests

Commonly recommended samples to detect ASFV, its nucleic acids and antigens include blood from live animals or tissues (especially spleen, kidney, tonsils, lymph nodes, liver, heart and lung) collected at necropsy. The spleen and lymph nodes usually contain the highest concentrations of virus, and viral DNA may persist longer in the spleen than other internal organs after death. Nucleic acids may also be detected in the bone marrow, which can be useful when other tissues from carcasses are not available or usable, and the intra-articular tissues of joints are sometimes tested in chronic cases. Most secretions and excretions are also likely to contain virus during the clinical stage in live animals, but there is little or no published information for using these samples in diagnostic testing, except in aggregate pen-based surveillance, such as ropes for saliva testing. Other types of samples, such as ear punches, have also been investigated for PCR-based surveillance of carcasses.

A blood sample from the dam has usually been recommended in cases of abortion, as ASFV was not thought to occur in the fetus. However, some fetuses were recently found to have at least low levels of viral DNA in tissues such as the tonsils, spleen, lymph nodes, liver and lung, as well as the placenta. ASFV was successfully isolated from the placenta, amniotic fluid and fetal blood after abortions caused by a low virulence isolate, though the fetal tissues showed little or no immunostaining for viral antigens. Both viral nucleic acids and antigens were detected in the brain and cerebrospinal fluid of neonatal piglets with congenital



tremors; however, the virus appeared to be absent from other tissues including the spleen, kidney and lymph nodes.

Clinical cases of African swine fever are often diagnosed with various PCR tests. Loop-mediated isothermal amplification (LAMP), recombinase polymerase amplification (RPA) and recombinase-aided amplification (RAA) assays to detect ASFV nucleic acids have also been published. ASFV antigens may be found in tissues or buffy coat samples with ELISAs or immunofluorescence. Rapid penside lateral flow devices for antigen detection have also been described, and may be available as commercial tests in some countries. Antigens are easiest to find in acute cases, and antigen-detection tests sometimes have relatively low sensitivity; thus, they are best employed as herd tests, in conjunction with other assays. A hemadsorption “autorosette” test was also used in the past to detect ASFV directly in peripheral blood leukocytes, but it has mostly been replaced by PCR.

If necessary, ASFV can be isolated in primary porcine cells including pig leukocyte or bone marrow cultures, porcine alveolar macrophages or blood monocyte cultures. Primary cells are complicated to work with, and reports of cell lines that might be able to replace these cells, such as MA-104 cells, have been published. Additional cell lines (e.g., COS-1 cells, Vero cells) can be used to propagate viruses that have been adapted to cell culture, but none of these cells is susceptible to all field strains, limiting their usefulness for diagnosis. Traditionally, ASFV-infected cells have been identified in cultures by their ability to induce hemadsorption of pig erythrocytes to their surfaces, but a few non-hemadsorbing isolates, including some that cause chronic disease, can be missed with this test. Virus-infected cells can also be detected by PCR or immunofluorescence, and PCR can be used to confirm the virus’s identity.

Pigs with acute disease often die before seroconverting; however antibodies to ASFV persist for long periods in animals that survive. The currently used serological assays include ELISAs, immunoblotting, indirect fluorescent antibody (IFA) and indirect immunoperoxidase (IPT) tests, and penside lateral flow assays to detect antibodies have been commercialized in some countries. The ELISA is prescribed for international trade, and is generally confirmed by immunoblotting, but IFA or IPT can also be used for confirmation.

## Treatment

There is no treatment for African swine fever, other than supportive care.

## Control

### Disease reporting

Veterinarians who encounter or suspect African swine fever should follow their national and/or local guidelines for disease reporting. In the U.S., state or federal veterinary authorities should be informed immediately.

### Prevention

Biosecurity measures, including quarantines of new animals, reduce the risk of introducing ASFV onto a farm. Separation of the herd from wild suids, their environments and carcasses (e.g., with double fencing), must also be considered in many areas. In the past, many ASFV-free countries used heat treatment to inactivate any viruses that might be present in pig swill. Due to the risk that ASFV and other viruses may not be completely inactivated (for example, if parts of the swill do not reach the target temperature), some nations have completely forbidden feeding swill to pigs. In areas where this is not feasible, some sources recommend boiling the swill for at least 30 minutes, with frequent or continuous stirring. Solid walls without cracks are considered the optimal building material to discourage the establishment of *Ornithodoros* ticks and facilitate control. Acaricides are generally ineffective where wooden, stone, earth or overlapping metal walls/ fences provide hiding places for these ticks.

Some areas have successfully eradicated African swine fever outbreaks in domesticated pigs with standard stamping out measures (e.g., slaughter of infected and in-contact animals, sanitation, disinfection, movement controls and quarantines), but more complex measures were needed in some regions. On the Iberian Peninsula, ASFV became established in wild boar and *Ornithodoros erraticus* ticks in the 1960s, and complete eradication took decades. Piggens with infected ticks were destroyed or isolated as part of this campaign, and minimum quarantines were much longer (e.g., 6 years rather than 40 days) where the tick vectors were thought to be involved in transmission. Because ASFV often spreads relatively slowly through a herd, some Asian countries where this virus has become common have recently implemented partial or selective culling of herds, rather than complete depopulation, as part of their control efforts. The success of this measure remains to be determined, but potential benefits include an incentive for early reporting of outbreaks, as well as reduced losses for farmers.

Current control measures in wild boar mainly focus on reducing their numbers by hunting and/or trapping, as higher population densities are thought to facilitate maintenance of the virus, and attempts to discourage the movement of infected animals with fencing or other means. In some countries, authorities also search for and remove wild boar carcasses. Campaigns to eliminate the virus from wild boar populations with these measures have often met with limited success, and appear to be more effective when there is a single geographical focus of infection. Biosecurity measures to reduce the risk of transporting ASFV during hunting (e.g., the use of leak-proof vessels to remove carcasses and store offal, limits on vehicles in infected areas, precautions for cleaning and disinfecting tools) have also been published. Eradication of ASFV from some wild reservoirs in Africa, such as warthogs, appears unlikely, but compartments where African swine fever is controlled and barriers (double fencing) prevent contact with wild reservoirs have been established in some regions where warthog-mediated introduction is a concern.

Vaccines have been difficult to develop against ASFV, which has a large and complex genome. Although two live attenuated vaccines have been licensed in Vietnam, pigs can shed the vaccine viruses, raising concerns about reversion to virulence or recombination with field strains. Some vaccinated animals also experienced health problems or died, and both vaccine viruses were recently shown to revert to virulence on serial passage in pigs. One resulted in mild clinical signs and prolonged viremia after 5 passages, and the other caused severe clinical signs and high mortality after 3-4 passages.

## Morbidity and Mortality

ASFV currently circulates in warthogs as well as domesticated pigs in parts of Africa, and in both pigs and wild boar in parts of Eurasia. While previous experiences in Europe suggested that this virus would eventually die out in wild boar if it was not reintroduced from pigs, this has not happened. Instead, it has spread slowly in boar populations and diversified, independently of its occurrence in farmed pigs. In Estonia, initial high mortality rates among wild boar, together with control measures, resulted in a significant decline in the numbers of these animals and the apparent disappearance of the virus; however, it continued to circulate at low levels (which were not detected at the time by surveillance) and the recovery of the boar population was accompanied by the resurgence of the virus.

The importance of wild boar in infecting domesticated pigs (or vice versa) may differ between regions. In some areas, especially those with large numbers of backyard pigs, most farms seem to become infected from other farms rather than wild suids. Outbreaks appear to be somewhat seasonal in both pigs and wild boar, though the reported patterns differ between pigs and boar, and between farms in different countries. Patterns of pig marketing, including changes in the demand for pork products, and other cultural factors are generally thought to account for the seasonality in domestic herds, though it remains possible that other factors, such as arthropod-mediated transmission, might also play a role. In wild boar, seasonality might be caused by changes in wild boar interactions (e.g., mating season), as well as longer virus survival under cold conditions; however, it is possible that the observed variations are just an artifact of human observations, for instance the increased likelihood of detecting dead boar carcasses or infected animals at times when more people are in the woods.

The morbidity and mortality rates for African swine fever can be influenced by the virulence of the isolate, the dose of the virus, and animal factors such as previous exposures and general health. Morbidity rates can approach 100% in some naive herds of domesticated pigs, while cumulative mortality ranges from < 5% with mild or attenuated viruses to 100% from highly virulent viruses. The mortality rate is usually around 30-70% in subacute cases. Less virulent isolates are more likely to kill pigs with concurrent diseases, pregnant or nursing sows, and young animals. Morbidity and mortality rates also tend to be higher

when ASFV is introduced into a new region, with an increased incidence of milder cases once it becomes endemic. Because ASFV sometimes takes days to weeks to spread through a herd, initial herd mortality rates may be low even when the case fatality rate is high.

Both domesticated pig and wild boar populations can maintain a variety of ASF isolates of varying virulence. Some viruses have been shown to cause more severe disease in wild boar than pigs, though it is not known whether this is true for all or most strains. Chronic African swine fever was first described during outbreaks on the Iberian Peninsula, leading to speculation that the viruses that cause this form might have originated from live attenuated vaccine strains tested at the time. However, it can also be caused by virulent viruses, including some recent Eurasian strains. Pigs that recover from African swine fever are often resistant to illness upon subsequent exposure, though they may become reinfected and shed the virus. However, this protection does not necessarily extend to all other strains. Certain populations of pigs in Africa seem to be relatively resistant to African swine fever, but the basis for this resistance is not known.

## Internet Resources

[CIRAD Pigdrop \(Pig Production in Developing Countries\)](#)

[Food and Agriculture Organization of the United Nations \(FAO\). Recognizing African Swine Fever. A Field Manual.](#)

[FAO. Updates on the ASF Situation Worldwide \(with links to African swine fever information\)](#)

[FAO: African Swine Fever: Detection and Diagnosis – A Manual for Veterinarians \(English version\)](#)

[The Merck Veterinary Manual](#)

[United States Animal Health Association. Foreign Animal Diseases](#)

[World Organization for Animal Health \(WOAH\)](#)

[WOAH Manual of Diagnostic Tests and Vaccines for Terrestrial Animals; WOAH Terrestrial Animal Health Code](#)

## Acknowledgements

This factsheet was written by Anna Rovid Spickler, DVM, PhD, Veterinary Specialist from the Center for Food Security and Public Health. The U.S. Department of Agriculture Animal and Plant Health Inspection Service (USDA APHIS) provided funding for this factsheet through a series of cooperative agreements related to the development of resources for initial accreditation training.

The following format can be used to cite this factsheet. Spickler, Anna Rovid. 2025. *African Swine Fever*. Retrieved from <http://www.cfsph.iastate.edu/DiseaseInfo/factsheets.php>.

## References

- Animal Health Australia. The National Animal Health Information System (NAHIS). African swine fever [online]. Available at: [http://www.brs.gov.au/usr-bin/aphb/ahsq?dislist=alpha.\\*](http://www.brs.gov.au/usr-bin/aphb/ahsq?dislist=alpha.*) Accessed 18 Oct 2001.
- Arias M, Jurado C, Gallardo C, Fernández-Pinero J, Sánchez-Vizcaíno JM. Gaps in African swine fever: Analysis and priorities. *Transbound Emerg Dis*. 2018;65 Suppl 1:235-47.
- Arzumanyan H, Hakobyan S, Avagyan H, Izmailyan R, Nersisyan N, Karalyan Z. Possibility of long-term survival of African swine fever virus in natural conditions. *Vet World*. 2021;14(4):854-9.
- Auer A, Cattoli G, Padungtod P, Lamien CE, Oh Y, Jayme S, Rozstalnyy A. Challenges in the application of African swine fever vaccines in Asia. *Animals (Basel)*. 2024;14(17):2473.
- Avagyan HR, Hakobyan SA, Avetisyan AS, Bayramyan NV, Hakobyan LH, Poghosyan AA, Abroyan LO, Baghdasaryan BV, Tsakanova GV, Sahakyan LV, Yeremyan AS, Karalyan ZA. The pattern of stability of African swine fever virus in leeches. *Vet Microbiol*. 2023;284:109835.
- Ayoade GO; Adeyemi IG. African swine fever: an overview. *Revue Elev Méd vét Pays Trop*. 2003;56:129-34.
- Balmoş OM, Supeanu A, Tamba P, Horváth C, Panait LC, Sándor AD, Cazan CD, Ungur A, Motiu M, Manita FA, Ancuceanu BC, Bărbuceanu F, Dhollander S, Mihalca AD. African swine fever virus load in hematophagous dipterans collected in outbreaks from Romania: risk factors and implications. *Transbound Emerg Dis*. 2023;2023:3548109.
- Beato MS, D'Errico F, Iscaro C, Petrini S, Giammarioli M, Feliziani F. Disinfectants against African swine fever: an updated review. *Viruses*. 2022;14(7):1384.
- Bellini S, Rutili D, Guberti V. Preventive measures aimed at minimizing the risk of African swine fever virus spread in pig farming systems. *Acta Vet Scand*. 2016;58(1):82.
- Beltrán-Alcrudo D, Arias M, Gallardo C, Kramer S, Penrith ML. African swine fever: detection and diagnosis – A manual for veterinarians. *FAO Animal Production and Health Manual No. 19*. Rome: Food and Agriculture Organization of the United Nations (FAO); 2017. Available at: <https://openknowledge.fao.org/bitstreams/d6d6ad7b-b4d7-4116-a95e-06826e8a25b0/download>. Accessed 18 Jun 2019.
- Blázquez E, Pujols J, Segalés J, Rodríguez F, Crenshaw J, Rodríguez C, Ródenas J, Polo J. Commercial feed containing porcine plasma spiked with African swine fever virus is not infective in pigs when administered for 14 consecutive days. *PLoS One*. 2020;15(7):e0235895.
- Blome S, Gabriel C, Beer M. Pathogenesis of African swine fever in domestic pigs and European wild boar. *Virus Res*. 2013;173(1):122-30.
- Boinas FS, Wilson AJ, Hutchings GH, Martins C, Dixon LJ. The persistence of African swine fever virus in field-infected *Ornithodoros erraticus* during the ASF endemic period in Portugal. *PLoS One*. 2011;6(5):e20383.
- Bonnet SI, Bouhsira E, De Regge N, Fite J, Etoré F, Garigliany MM, Jori F, Lempereur L, Le Potier MF, Quillery E, Saegerman C, Vergne T, Vial L. Putative role of arthropod vectors in African swine fever virus transmission in relation to their bio-ecological properties. *Viruses*. 2020;12(7):778.
- Chandana MS, Nair SS, Chaturvedi VK, Abhishek, Pal S, Charan MSS, Balaji S, Saini S, Vasavi K, Deepa P. Recent progress and major gaps in the vaccine development for African swine fever. *Braz J Microbiol*. 2024;55(1):997-1010.
- Chen S, Wang T, Luo R, Lu Z, Lan J, Sun Y, Fu Q, Qiu HJ. Genetic variations of African swine fever virus: major challenges and prospects. *Viruses*. 2024;16(6):913.
- Chenais E, Ahlberg V, Andersson K, Banihashem F, Björk L, et al. First outbreak of African swine fever in Sweden: local epidemiology, surveillance, and eradication strategies. *Transbound Emerg Dis*. 2024;2024:6071781.
- Chenais E, Depner K, Guberti V, Dietze K, Viltrop A, Ståhl K. Epidemiological considerations on African swine fever in Europe 2014-2018. *Porcine Health Manag*. 2019;5:6.
- Costard S, Mur L, Lubroth J, Sanchez-Vizcaino JM, Pfeiffer DU. Epidemiology of African swine fever virus. *Virus Res*. 2013;173(1):191-7.
- Costard S, Wieland B, de Glanville W, Jori F, Rowlands R, Vosloo W, Roger F, Pfeiffer DU, Dixon LK. African swine fever: how can global spread be prevented? *Philos Trans R Soc Lond B Biol Sci*. 2009;364(1530):2683-96.
- Cubillos C, Gómez-Sebastian S, Moreno N, Nuñez MC, Mulumba-Mfumu LK, Quembo CJ, Heath L, Etter EM, Jori F, Escribano JM, Blanco E. African swine fever virus serodiagnosis: a general review with a focus on the analyses of African serum samples. *Virus Res*. 2013;173(1):159-67.
- Dardiri AH, Yedloutschnig RJ, Taylor WD. Clinical and serologic response of American white-collared peccaries to African swine fever, foot-and-mouth disease, vesicular stomatitis, vesicular exanthema of swine, hog cholera, and rinderpest viruses. *Proc Annu Meet U S Anim Health Assoc*. 1969;73:437-52.
- Davies K, Goatley LC, Guinat C, Netherton CL, Gubbins S, Dixon LK, Reis AL. Survival of African swine fever virus in excretions from pigs experimentally infected with the Georgia 2007/1 isolate. *Transbound Emerg Dis*. 2017;64(2):425-31.
- de Carvalho Ferreira HC, Tudela Zúquete S, Wijnveld M, Weesendorp E, Jongejan F, Stegeman A, Loeffen WL. No evidence of African swine fever virus replication in hard ticks. *Ticks Tick Borne Dis*. 2014;5(5):582-9.
- de Carvalho Ferreira HC, Weesendorp E, Quak S, Stegeman JA, Loeffen WL. Quantification of airborne African swine fever virus after experimental infection. *Vet Microbiol*. 2013;165(3-4):243-51.
- de Carvalho Ferreira HC, Weesendorp E, Quak S, Stegeman JA, Loeffen WL. Suitability of faeces and tissue samples as a basis for non-invasive sampling for African swine fever in wild boar. *Vet Microbiol*. 2014;172(3-4):449-54.
- De Lorenzi G, Borella L, Alborali GL, Prodanov-Radulović J, Štukelj M, Bellini S. African swine fever: A review of cleaning and disinfection procedures in commercial pig holdings. *Res Vet Sci*. 2020;132:262-7.
- Deutschmann P, Pikalo J, Beer M, Blome S. Lateral flow assays for the detection of African swine fever virus antigen are not fit for field diagnosis of wild boar carcasses. *Transbound Emerg Dis*. 2022;69(4):2344-8.
- Diaz AV, Netherton CL, Dixon LK, Wilson AJ. African swine fever virus strain Georgia 2007/1 in *Ornithodoros erraticus* ticks. *Emerg Infect Dis*. 2012;18(6):1026-8.

- Dixon LK, Sun H, Roberts H. African swine fever. *Antiviral Res.* 2019;165:34-41.
- Duc HM, Ngan PH, Son HM, Lan NT, Van Hung L, Ha CTT, Hoa NT, Lam TQ, Van Thang N, Flory GA, Hutchinson M. The use of composting for the disposal of African swine fever virus-infected swine carcasses. *Transbound Emerg Dis.* 2022;69(5):e3036-44.
- Eblé PL, Hagenaars TJ, Weesendorp E, Quak S, Moonen-Leusen HW, Loeffen WLA. Transmission of African swine fever virus via carrier (survivor) pigs does occur. *Vet Microbiol.* 2019;237:108345.
- EFSA Panel on Animal Health and Welfare (EFSA AHAW Panel); Nielsen SS, Alvarez J, Bicout DJ, Calistri P, Depner K, et al. Scientific Opinion on the assessment of the control measures of the category A diseases of Animal Health Law: African Swine Fever. *EFSA J.* 2021;19(1):e06402.
- Endris RG, Hess WR. Experimental transmission of African swine fever virus by the soft tick *Ornithodoros (Pavlovskyella) maroccanus* (Acari: Ixodoidea: Argasidae). *J Med Entomol.* 1992;29:652-6.
- European Food Safety Authority (EFSA); Ståhl K, Boklund AE, Podgórski T, Vergne T, Abrahantes JC, Cattaneo E, Papanikolaou A, Mur L. Epidemiological analysis of African swine fever in the European Union during 2023. *EFSA J.* 2024;22(5):e8809.
- European Food Safety Authority (EFSA); Ståhl K, Boklund AE, Podgórski T, Vergne T, Aminalragia-Giamini R, Abrahantes JC, Papaleo S, Mur L. Epidemiological analysis of African swine fever in the European Union during 2024. *EFSA J.* 2025; 19;23(5):e9436.
- EFSA Panel on Animal Health and Welfare (AHAW); Nielsen SS, Alvarez J, Bicout DJ, Calistri P, Canali E, et al. African swine fever and outdoor farming of pigs. *EFSA J.* 2021;19(6):e06639.
- Fischer M, Hühner J, Blome S, Conraths FJ, Probst C. Stability of African swine fever virus in carcasses of domestic pigs and wild boar experimentally infected with the ASFV "Estonia 2014" isolate. *Viruses.* 2020;12(10):1118.
- Food and Agriculture Organization of the United Nations (FAO). Recognizing African swine fever. A field manual [online]. FAO Animal Health Manual No. 9. Rome: FAO; 2000. Available at: <http://www.fao.org/docrep/004/X8060E/X8060E00.HTM>. Accessed 4 Dec 2006.
- Forth JH, Calvelage S, Fischer M, Hellert J, Sehl-Ewert J, et al. African swine fever virus - variants on the rise. *Emerg Microbes Infect.* 2023;12(1):2146537.
- Friedrichs V., Reicks D., Hasenfuss T., Gerstenkorn E., Zimmermann J., Nelson E., Carrau T., Deutschmann P., Sehl-Ewert J., Roszyk H., et al. Artificial insemination as alternative transmission route for African swine fever virus. *Pathogens.* 2022;11:1539.
- Frost L, Tully M, Dixon L, Hicks HM, Bennett J, Stokes I, Marsella L, Gubbins S, Batten C. Evaluation of the efficacy of commercial disinfectants against African swine fever virus. *Pathogens.* 2023;12(7):855.
- Gallardo C, Fernández-Pinero J, Arias M. African swine fever (ASF) diagnosis, an essential tool in the epidemiological investigation. *Virus Res.* 2019;271:197676.
- Gallardo C, Fernández-Pinero J, Pelayo V, Gazeau I, Markowska-Daniel I, Pridotkas G, Nieto R, Fernández-Pacheco P, Bokhan S, Nevolko O, Drozhzhe Z, Pérez C, Soler A, Kolvasov D, Arias M. Genetic variation among African swine fever genotype II viruses, eastern and central Europe. *Emerg Infect Dis.* 2014;20(9):1544-7.
- Gallardo C, Nurmoja I, Soler A, Delicado V, Simón A, Martín E, Pérez C, Nieto R, Arias M. Evolution in Europe of African swine fever genotype II viruses from highly to moderately virulent. *Vet Microbiol.* 2018;219:70-9.
- Gallardo C, Soler A, Nieto R, Sánchez MA, Martins C, Pelayo V, Carrascosa A, Revilla Y, Simón A, Briones V, Sánchez-Vizcaíno JM, Arias M. Experimental transmission of African swine fever (ASF) low virulent isolate NH/P68 by surviving pigs. *Transbound Emerg Dis.* 2015;62(6):612-22.
- Gallardo C, Soler A, Rodze I, Nieto R, Cano-Gómez C, Fernández-Pinero J, Arias M. Attenuated and non-haemadsorbing (non-HAD) genotype II African swine fever virus (ASFV) isolated in Europe, Latvia 2017. *Transbound Emerg Dis.* 2019;66(3):1399-404.
- Gavriel-Widén D, Gortázar C, Ståhl K, Neimanis AS, Rossi S, Hårdav Segerstad C, Kuiken T. African swine fever in wild boar in Europe: a notable challenge. *Vet Rec.* 2015;176(8):199-200.
- Giammarioli M, Gallardo C, Oggiano A, Iscaro C, Nieto R, Pellegrini C, Dei Giudici S, Arias M, De Mia GM. Genetic characterisation of African swine fever viruses from recent and historical outbreaks in Sardinia (1978-2009). *Virus Genes.* 2011;42(3):377-87.
- Glšić D, Milićević V, Veljović L, Milovanović B, Kureljušić B, Đorđević I, Anđelković K, Petković J, Đaćić M. Patterns of ASFV transmission in domestic pigs in Serbia. *Pathogens.* 2023;12(1):149.
- Guinat C, Gogin A, Blome S, Keil G, Pollin R, Pfeiffer DU, Dixon L. Transmission routes of African swine fever virus to domestic pigs: current knowledge and future research directions. *Vet Rec.* 2016;178(11):262-7.
- Guinat C, Reis AL, Netherton CL, Goatley L, Pfeiffer DU, Dixon L. Dynamics of African swine fever virus shedding and excretion in domestic pigs infected by intramuscular inoculation and contact transmission. *Vet Res.* 2014;45:93.
- Hakobyan SA, Ross PA, Bayramyan NV, Poghosyan AA, Avetisyan AS, Avagyan HR, Hakobyan LH, Abroyan LO, Harutyunova LJ, Karalyan ZA. Experimental models of ecological niches for African swine fever virus. *Vet Microbiol.* 2022;266:109365.
- Herm R, Kirik H, Vilem A, Zani L, Forth JH, Müller A, Michelitsch A, Wernike K, Werner D, Tummeleht L, Kampen H, Viltrop A. No evidence for African swine fever virus DNA in haematophagous arthropods collected at wild boar baiting sites in Estonia. *Transbound Emerg Dis.* 2021;68(5):2696-702.
- Hess WR, Endris RG, Lousa A, Caiado JM. Clearance of African swine fever virus from infected tick (Acari) colonies. *J Med Entomol.* 1989;26(4):314-7.
- Iglesias I, Rodríguez A, Feliziani F, Rolesu S, de la Torre A. Spatio-temporal analysis of African swine fever in Sardinia (2012–2014): trends in domestic pigs and wild boar. *Transbound Emerg Dis.* 2017;64:656-62.



- Igolkin A, Mazloun A, Zinyakov N, Chernyshev R, Schalkwyk AV, Shotin A, Lavrentiev I, Gruzdev K, Chvala I. Detection of the first recombinant African swine fever virus (genotypes I and II) in domestic pigs in Russia. *Mol Biol Rep.* 2024;51(1):1011.
- Ito S, Kawaguchi N, Bosch J, Aguilar-Vega C, Sánchez-Vizcaino JM. What can we learn from the five-year African swine fever epidemic in Asia? *Front Vet Sci.* 2023;10:1273417.
- Jo YS, Gortázar C. African swine fever in wild boar: Assessing interventions in South Korea. *Transbound Emerg Dis.* 2021;68(5):2878-89.
- Jori F, Bastos AD. Role of wild suids in the epidemiology of African swine fever. *Ecohealth.* 2009;6(2):296-310.
- Jori F, Bastos A, Boinas F, Van Heerden J, Heath L, Jourdan-Pineau H, Martinez-Lopez B, Pereira de Oliveira R, Pollet T, Quembo C, Rea K, Simulundu E, Taraveau F, Penrith ML. An updated review of *Ornithodoros* ticks as reservoirs of African swine fever in Sub-Saharan Africa and Madagascar. *Pathogens.* 2023;12(3):469.
- Juszkiewicz M, Walczak M, Woźniakowski G. Characteristics of selected active substances used in disinfectants and their virucidal activity against ASFV. *J Vet Res.* 2019;63(1):17-25.
- Juszkiewicz M, Walczak M, Woźniakowski G, Pejsak Z, Podgórska K. The influence of the temperature on effectiveness of selected disinfectants against African swine fever virus (ASFV). *Viruses.* 2025;17(2):156.
- Kleiboeker SB. African swine fever. In: *Foreign animal diseases.* Richmond, VA: United States Animal Health Association; 2008. p. 111-6.
- Kleiboeker SB. Swine fever: classical swine fever and African swine fever. *Vet Clin North Am Food Anim Pract.* 2002;18:431-51.
- Krug PW, Larson CR, Eslami AC, Rodriguez LL. Disinfection of foot-and-mouth disease and African swine fever viruses with citric acid and sodium hypochlorite on birch wood carriers. *Vet Microbiol.* 2012;156(1-2):96-101.
- Krug PW, Lee LJ, Eslami AC, Larson CR, Rodriguez L. Chemical disinfection of high-consequence transboundary animal disease viruses on nonporous surfaces. *Biologicals.* 2011;39(4):231-5.
- Laddomada A, Rolesu S, Loi F, Cappai S, Oggiano A et al. Surveillance and control of African swine fever in free-ranging pigs in Sardinia. *Transbound Emerg Dis.* 2019;66(3):1114-9.
- Le VP, Jeong DG, Yoon SW, Kwon HM, Trinh TBN, et al. Outbreak of African swine fever, Vietnam, 2019. *Emerg Infect Dis.* 2019;25(7):1433-5.
- Lee K, Vu TTH, Yeom M, Nguyen VD, Than TT, Nguyen VT, Jeong DG, Ambagala A, Le VP, Song D. Molecular characterization of emerging recombinant African swine fever virus of genotype I and II in Vietnam, 2023. *Emerg Microbes Infect.* 2024;13(1):2404156.
- Li L, Ren Z, Wang Q, Ge S, Liu Y, et al. Infection of African swine fever in wild boar, China, 2018. *Transbound Emerg Dis.* 2019;66(3):1395-8.
- Li X, Hu Z, Fan M, Tian X, Wu W, Gao W, Bian L, Jiang X. Evidence of aerosol transmission of African swine fever virus between two piggeries under field conditions: a case study. *Front Vet Sci.* 2023;10:1201503.
- Linden A, Licoppe A, Volpe R, Paternostre J, Lesenfans C, Cassart D, Garigliany M, Tignon M, van den Berg T, Desmecht D, Cay AB. Summer 2018: African swine fever virus hits north-western Europe. *Transbound Emerg Dis.* 2019;66(1):54-5.
- Lohse L, Nielsen J, Uttenthal Å, Olesen AS, Strandbygaard B, Rasmussen TB, Belsham GJ, Bøtner A. Experimental infections of pigs with African swine fever virus (genotype II); studies in young animals and pregnant sows. *Viruses.* 2022;14(7):1387.
- Loundras EA, Netherton CL, Flannery J, Bowes MJ, Dixon L, Batten C. The effect of temperature on the stability of African swine fever virus BA71V isolate in environmental water samples. *Pathogens.* 2023;12(8):1022.
- Mazur-Panasiuk N, Woźniakowski G. Natural inactivation of African swine fever virus in tissues: Influence of temperature and environmental conditions on virus survival. *Vet Microbiol.* 2020;242:108609.
- Mebus CA, Arias M, Pineda JM, Tapiador J, House C, Sanchez-Vizcaino JM. Survival of several porcine viruses in Spanish dry-cured meat products. *Food Chem* 1997;59:555-9.
- Mebus CA, Dardiri AH. Additional characteristics of disease caused by the African swine fever viruses isolated from Brazil and the Dominican Republic. *Proc Ann Meet US Anim Health Ass.* 1979;82:227-39.
- More S, Miranda MA, Bicout D, Botner A, Butterworth A, et. al.; EFSA Panel on Animal Health and Welfare. African swine fever in wild boar. *ESFA J.* 2018;16(7):5344.
- Mulumba-Mfumum LK, Saegerman C, Dixon LK, Madimba KC, Kazadi E, Mukalakata NT, Oura CAL, Chenais E, Masembe C, Ståhl K, Thiry E, Penrith ML. African swine fever: Update on eastern, central and southern Africa. *Transbound Emerg Dis.* 2019 66(4):1462-80.
- Mur L, Boadella M, Martínez-López B, Gallardo C, Gortazar C, Sánchez-Vizcaino JM. Monitoring of African swine fever in the wild boar population of the most recent endemic area of Spain. *Transbound Emerg Dis.* 2012;59(6):526-31.
- Mushagalusa CA, Etter E, Penrith ML. Review of African swine fever outbreaks history in South Africa: From 1926 to 2018. *Onderstepoort J Vet Res.* 2021;88(1):e1-10.
- Niederwerder MC, Khanal P, Foland T, Constance LA, Stoian AMM, Deavours A, Haase K, Cino-Ozuna AG. Stability of African swine fever virus in feed during environmental storage. *Transbound Emerg Dis.* 2022;69(6):3216-24.
- Nga BTT, Padungtod P, Depner K, Chuong VD, Duy DT, Anh ND, Dietze K(3) Implications of partial culling on African swine fever control effectiveness in Vietnam. *Front Vet Sci.* 2022;9:957918.
- Nguyen HN, Nguyen QT, Nguyen BTP, Tran THA, Do DT, Hoang HT. Detection of African swine fever virus in neonatal piglets with congenital tremors. *Arch Virol.* 2022;167(4):1131-5.
- Niederwerder MC, Stoian AMM, Rowland RRR, Dritz SS, Petrovan V, Constance LA, Gebhardt JT, Olcha M, Jones CK, Woodworth JC, Fang Y, Liang J, Hefley TJ. Infectious dose of African swine fever virus when consumed naturally in liquid or feed. *Emerg Infect Dis.* 2019;25(5):891-7.

- Nuanualsuwan S, Songkasupa T, Boonpornprasert P, Suwankitwat N, Lohlamoh W, Nuengjamnong C. Thermal inactivation of African swine fever virus in swill. *Front Vet Sci*. 2022;9:906064.
- Nurmoja I, Mõtus K, Kristian M, Niine T, Schulz K, Depner K, Viltrop A. Epidemiological analysis of the 2015-2017 African swine fever outbreaks in Estonia. *Prev Vet Med*. 2020;181:104556.
- Oh T, Nguyen TM, Ngo TTN, Thinh D, Nguyen TTP, Do LD, Do DT. Long-term follow-up of convalescent pigs and their offspring after an outbreak of acute African swine fever in Vietnam. *Transbound Emerg Dis*. 2021;68(6):3194-9.
- Olesen AS, Hansen MF, Rasmussen TB, Belsham GJ, Bødker R, Bøtner A. Survival and localization of African swine fever virus in stable flies (*Stomoxys calcitrans*) after feeding on viremic blood using a membrane feeder. *Vet Microbiol*. 2018;222:25-9.
- Olesen AS, Lohse L, Accensi F, Goldswain H, Belsham GJ, Bøtner A, Netherton CL, Dixon LK, Portugal R. Inefficient transmission of African swine fever virus to sentinel pigs from an environment contaminated by ASFV-infected pigs under experimental conditions. *Transbound Emerg Dis*. 2024;2024:8863641.
- Olesen AS, Lohse L, Boklund A, Halasa T, Belsham GJ, Rasmussen TB, Bøtner A. Short time window for transmissibility of African swine fever virus from a contaminated environment. *Transbound Emerg Dis*. 2018;65(4):1024-32.
- Olesen AS, Lohse L, Boklund A, Halasa T, Gallardo C, Pejsak Z, Belsham GJ, Rasmussen TB, Bøtner A. Transmission of African swine fever virus from infected pigs by direct contact and aerosol routes. *Vet Microbiol*. 2017;211:92-102.
- Olesen AS, Lohse L, Hansen MF, Boklund A, Halasa T, Belsham GJ, Rasmussen TB, Bøtner A, Bødker R. Infection of pigs with African swine fever virus via ingestion of stable flies (*Stomoxys calcitrans*). *Transbound Emerg Dis*. 2018;65(5):1152-7.
- Olesen AS, Stelder JJ, Tjørnehøj K, Johnston CM, Lohse L, Kjær LJ, Boklund AE, Bøtner A, Belsham GJ, Bødker R, Rasmussen TB. Detection of African swine fever virus and blood meals of porcine origin in hematophagous insects collected adjacent to a high-biosecurity pig farm in Lithuania; a smoking Ggn? *Viruses*. 2023;15(6):1255.
- Oura C. Overview of African swine fever. In: Kahn CM, Line S, Aiello SE, editors. *The Merck veterinary manual*. 10th ed. Whitehouse Station, NJ: Merck and Co; 2013. Available at: [http://www.merckvetmanual.com/mvm/generalized\\_condition/s/african\\_swine\\_fever/overview\\_of\\_african\\_swine\\_fever.html](http://www.merckvetmanual.com/mvm/generalized_condition/s/african_swine_fever/overview_of_african_swine_fever.html). Accessed 15 Oct 2015.
- Oura CA, Edwards L, Batten CA. Virological diagnosis of African swine fever--comparative study of available tests. *Virus Res*. 2013;173(1):150-8.
- Pan L, Luo R, Wang T, Qi M, Wang B, Sun M, Luo Y, Ji C, Sun Y, Qiu HJ. Efficient inactivation of African swine fever virus by a highly complexed iodine. *Vet Microbiol*. 2021;263:109245.
- Penrith ML, Bastos AD, Etter EMC, Beltrán-Alcruído D. Epidemiology of African swine fever in Africa today: Sylvatic cycle versus socio-economic imperatives. *Transbound Emerg Dis*. 2019;66(2):672-86.
- Penrith ML, Kivaria FM. One hundred years of African swine fever in Africa: Where have we been, where are we now, where are we going? *Transbound Emerg Dis*. 2022;69(5):e1179-1200.
- Penrith ML, van Emmenes J, Hakizimana JN, Heath L, Kabuuka T, Misinzio G, Odoom T, Wade A, Zerbo HL, Luka PD. African swine fever diagnosis in Africa: challenges and opportunities. *Pathogens*. 2024;13(4):296.
- Penrith ML, Van Heerden J, Heath L, Abworo EO, Bastos ADS. Review of the pig-adapted African swine fever viruses in and outside Africa. *Pathogens*. 2022;11(10):1190.
- Pereira de Oliveira R, Hutet E, Duhayon M, Guionnet JM, Paboeuf F, Vial L, Le Potier MF. Successful infection of domestic pigs by ingestion of the European soft tick *O. erraticus* that fed on African swine fever virus infected pig. *Viruses*. 2020;12(3):300.
- Pereira de Oliveira R, Hutet E, Lancelot R, Paboeuf F, Duhayon M, Boinas F, Pérez de León AA, Filatov S, Le Potier MF, Vial L. Differential vector competence of *Ornithodoros* soft ticks for African swine fever virus: What if it involves more than just crossing organic barriers in ticks? *Parasit Vectors*. 2020;13(1):618.
- Pereira de Oliveira R, Hutet E, Paboeuf F, Duhayon M, Boinas F, Perez de Leon A, Filatov S, Vial L, Le Potier MF. Comparative vector competence of the Afrotropical soft tick *Ornithodoros moubata* and Palearctic species, *O. erraticus* and *O. verrucosus*, for African swine fever virus strains circulating in Eurasia. *PLoS One*. 2019;14(11):e0225657.
- Petrini S, Feliziani F, Casciari C, Giammarioli M, Torresi C, De Mia GM. Survival of African swine fever virus (ASFV) in various traditional Italian dry-cured meat products. *Prev Vet Med*. 2019;162:126-30.
- Petrov A, Forth JH, Zani L, Beer M, Blome S. No evidence for long-term carrier status of pigs after African swine fever virus infection. *Transbound Emerg Dis*. 2018;65(5):1318-28.
- Pietschmann J, Guinat C, Beer M, Pronin V, Tauscher K, Petrov A, Keil G, Blome S. Course and transmission characteristics of oral low-dose infection of domestic pigs and European wild boar with a Caucasian African swine fever virus isolate. *Arch Virol*. 2015;160(7):1657-67.
- Pikalo J, Deutschmann P, Fischer M, Roszyk H, Beer M, Blome S. African swine fever laboratory diagnosis-lessons learned from recent animal trials. *Pathogens*. 2021;10(2):177.
- Pikalo J, Zani L, Hühr J, Beer M, Blome S. Pathogenesis of African swine fever in domestic pigs and European wild boar - lessons learned from recent animal trials. *Virus Res*. 2019;271:197614.
- Podgórski T, Śmietanka K. Do wild boar movements drive the spread of African swine fever? *Transbound Emerg Dis*. 2018;65(6):1588-96.
- Prodelalova J, Kavanova L, Salat J, Moutelikova R, Kobzova S, Krasna M, Vasickova P, Simek B, Vaclavek P. Experimental evidence of the long-term survival of infective African swine fever virus strain Ba71V in soil under different conditions. *Pathogens*. 2022;11(6):648.
- Radulovic E, Mehinagic K, Wüthrich T, Hilty M, Posthaus H, Summerfield A, Ruggli N, Benarafa C. The baseline immunological and hygienic status of pigs impact disease severity of African swine fever. *PLoS Pathog*. 2022;18(8):e1010522.

- Rahimi P, Sohrabi A, Ashraffihelan J, Edalat R, Alamdari M, Masoudi M, Mostofi S, Azadmanesh K. Emergence of African swine fever virus, northwestern Iran. *Emerg Infect Dis*. 2010;16(12):1946-8.
- Rai A, Pruitt S, Ramirez-Medina E, Vuono EA, Silva E, Velazquez-Salinas L, Carrillo C, Borca MV, Gladue DP. Identification of a continuously stable and commercially available cell line for the identification of infectious African swine fever virus in clinical samples. *Viruses*. 2020;12(8):820.
- Ramirez-Medina E, O'Donnell V, Silva E, Espinoza N, Velazquez-Salinas L, Moran K, Daite DA, Barrette R, Faburay B, Holland R, Gladue DP, Borca MV. Experimental infection of domestic pigs with an African swine fever virus field strain isolated in 2021 from the Dominican Republic. *Viruses*. 2022;14(5):1090.
- Ravaomanana J, Michaud V, Jori F, Andriatsimahavandy A, Roger F, Albina E. First detection of African swine fever Virus in *Ornithodoros porcinus* in Madagascar and new insights into tick distribution and taxonomy. *Parasit Vectors*. 2010;3:115.
- Ribeiro R, Otte J, Madeira S, Hutchings GH, Boinas F. Study of factors involved in the dynamics of infection in ticks. Experimental infection of *Ornithodoros erraticus sensu stricto* with two Portuguese African swine fever virus strains. *PLoS One*. 2015;10(9):e0137718.
- Rodríguez-Bertos A, Cadenas-Fernández E, Rebolada-Merino A, Porras-González N, Mayoral-Alegre FJ, Barreno L, Kosowska A, Tomé-Sánchez I, Barasona JA, Sánchez-Vizcaino JM. Clinical course and gross pathological findings in wild boar infected with a highly virulent strain of African swine fever virus genotype II. *Pathogens*. 2020;9(9):688.
- Rogoll L, Güttner AK, Schulz K, Bergmann H, Staubach C, Conraths FJ, Sauter-Louis C. Seasonal occurrence of African swine fever in wild boar and domestic pigs in EU member states. *Viruses*. 2023;15(9):1955.
- Sai Balaji KG, Karikalalan M, Saravanan S, Mahajan S, Rajeshkumar K, Vaseem M, Nautiyal S, Subash A, Sharma K, Biswas SK, Nandi S, Pawde AM, Singh KP, Sharma GK. High mortality in free-ranging wild boars associated with African swine fever virus in India. *Arch Virol*. 2024;169(7):137.
- Salguero FJ. Comparative pathology and pathogenesis of African swine fever infection in swine. *Front Vet Sci*. 2020;7:282.
- Sánchez-Cordón PJ, Floyd T, Hicks D, Croke HR, McCleary S, McCarthy RR, Strong R, Dixon LK, Neimanis A, Wikström-Lassa E, Gavier-Widén D, Núñez A. Evaluation of lesions and viral antigen distribution in domestic pigs inoculated intranasally with African swine fever virus Ken05/Tk1 (genotype X). *Pathogens*. 2021;10(6):768.
- Sánchez-Cordón PJ, Lean FZX, Batten C, Steinbach F, Neimanis A, Le Potier MF, Wikström-Lassa E, Wynne F, Strong R, McCleary S, Croke H, Gavier-Widén D, Núñez A. Comparative evaluation of disease dynamics in wild boar and domestic pigs experimentally inoculated intranasally with the European highly virulent African swine fever virus genotype II strain "Armenia 2007." *Vet Res*. 2024;55(1):89.
- Sánchez-Cordón PJ, Montoya M, Reis AL, Dixon LK. African swine fever: A re-emerging viral disease threatening the global pig industry. *Vet J*. 2018;233:41-8.
- Sanchez-Vizcaino JM. African swine fever from Kenya to five continents: the role of wild boar. *Rev Sci Tech*. 2024;Special Edition:53-7.
- Sánchez-Vizcaino JM, Mur L, Bastos AD, Penrith ML. New insights into the role of ticks in African swine fever epidemiology. *Rev Sci Tech*. 2015;34(2):503-11.
- Sánchez-Vizcaino JM, Mur L, Gomez-Villamandos JC, Carrasco L. An update on the epidemiology and pathology of African swine fever. *J Comp Pathol*. 2015;152(1):9-21.
- Sánchez-Vizcaino JM, Mur L, Martínez-López B. African swine fever: an epidemiological update. *Transbound Emerg Dis*. 2012;59 Suppl 1:27-35.
- Sargsyan MA, Voskanyan HE, Karalova EM, Hakobyan LH, Karalyan ZA. Third wave of African swine fever infection in Armenia: Virus demonstrates the reduction of pathogenicity. *Vet World*. 2018;11(1):5-9.
- Sastre P, Gallardo C, Monedero A, Ruiz T, Arias M, Sanz A, Rueda P. Development of a novel lateral flow assay for detection of African swine fever in blood. *BMC Vet Res*. 2016;12:206.
- Schambow RA, Hussain S, Antognoli MC, Kreindel S, Reyes R, Perez AM. Epidemiological assessment of African swine fever spread in the Dominican Republic. *Pathogens*. 2023;12(12):1414.
- Schulz K, Olševskis E, Staubach C, Lamberg K, Seržants M, Cvetkova S, Conraths FJ, Sauter-Louis C. Epidemiological evaluation of Latvian control measures for African swine fever in wild boar on the basis of surveillance data. *Sci Rep*. 2019;9(1):4189.
- Schulz K, Olševskis E, Viltrop A, Masiulis M, Staubach C, Nurmoja I, Lamberg K, Seržants M, Malakauskas A, Conraths FJ, Sauter-Louis C. Eight years of African swine fever in the Baltic states: epidemiological reflections. *Pathogens*. 2022;11(6):711.
- Schulz K, Schulz J, Staubach C, Blome S, Nurmoja I, Conraths FJ, Sauter-Louis C, Viltrop A. African swine fever re-emerging in Estonia: the role of seropositive wild boar from an epidemiological perspective. *Viruses*. 2021;13(11):2121.
- Schulz K, Staubach C, Blome S, Viltrop A, Nurmoja I, Conraths FJ, Sauter-Louis C. Analysis of Estonian surveillance in wild boar suggests a decline in the incidence of African swine fever. *Sci Rep*. 2019;9(1):8490.
- Sehl J, Pikalo J, Schäfer A, Franzke K, Pannhorst K, Elnagar A, Blohm U, Blome S, Breithaupt A. Comparative pathology of domestic pigs and wild boar infected with the moderately virulent African swine fever virus strain "Estonia 2014." *Pathogens*. 2020;9(8):662.
- Sehl-Ewert J, Deutschmann P, Breithaupt A, Blome S. Pathology of African swine fever in wild boar carcasses naturally infected with German virus variants. *Pathogens*. 2022;11(11):1386.
- Sehl-Ewert J, Friedrichs V, Carrau T, Deutschmann P, Blome S. Pathology of African swine fever in reproductive organs of mature breeding boars. *Viruses*. 2023;15(3):729.
- Shirai J, Kanno T, Tsuchiya Y, Mitsubayashi S, Seki R. Effects of chlorine, iodine, and quaternary ammonium compound disinfectants on several exotic disease viruses. *J Vet Med Sci*. 2000;62:85-92.

- Shurson GC, Palowski A, van de Ligt JLG, Schroeder DC, Balestreri C, Urriola PE, Sampedro F. New perspectives for evaluating relative risks of African swine fever virus contamination in global feed ingredient supply chains. *Transbound Emerg Dis.* 2022;69(1):31-56.
- Simulundu E, Chambaro HM, Sinkala Y, Kajihara M, Ogawa H, et al. Co-circulation of multiple genotypes of African swine fever viruses among domestic pigs in Zambia (2013-2015). *Transbound Emerg Dis.* 2018;65(1):114-22.
- Śmietanka K, Woźniakowski G, Kozak E, Niemczuk K, Frączyk M, Bocian Ł, Kowalczyk A, Pejsak Z. African swine fever epidemic, Poland, 2014-2015. *Emerg Infect Dis.* 2016;22(7):1201-7.
- Ståhl K, Sternberg-Lewerin S, Blome S, Viltrop A, Penrith ML, Chenais E. Lack of evidence for long term carriers of African swine fever virus - a systematic review. *Virus Res.* 2019;272:197725.
- Sun E, Huang L, Zhang X, Zhang J, Shen D, et al. Genotype I African swine fever viruses emerged in domestic pigs in China and caused chronic infection. *Emerg Microbes Infect.* 2021;10(1):2183-93.
- Torresi C, Biccheri R, Cammà C, Gallardo C, Marcacci M, et al. Genome-wide approach identifies natural large-fragment deletion in ASFV strains circulating in Italy during 2023. *Pathogens.* 2025;14(1):51.
- Urbano AC, Ferreira F. African swine fever control and prevention: an update on vaccine development. *Emerg Microbes Infect.* 2022;11(1):2021-33.
- van den Born E, Olasz F, Mészáros I, Göltl E, Oláh B, Joshi J, van Kilsdonk E, Segers R, Zádori Z. African swine fever virus vaccine strain Asfv-G-ΔI177I reverts to virulence and negatively affects reproductive performance. *NPJ Vaccines.* 2025;10(1):46.
- Vasić A, Milovanović B, Glišić D, Kavran M, Kureljušić J, Živulj A, Kureljušić B, Milićević V. The transmission routes of African swine fever during an outbreak in Serbia July-August 2023: African swine fever virus detections in environmental samples and insects. *Front Vet Sci.* 2024;11:1467273.
- Vial L, Wieland B, Jori F, Etter E, Dixon L, Roger F. African swine fever virus DNA in soft ticks, Senegal. *Emerg Infect Dis.* 2007;13(12):1928-31.
- Vu HD, Luong HQ, Lai HTL, Nguyen HT, Pham TH, Truong LQ, Nguyen GV, Vu HLX. Evaluation of the diagnostic sensitivity and specificity of two pen-side tests for detecting African swine fever virus in experimentally infected pigs. *Arch Virol.* 2024;169(8):170.
- Walczak M, Szymankiewicz K, Rodriguez F, Argilaguet J, Gavrilov B, Żmudzki J, Kochanowski M, Juskiewicz M, Szczotka-Bochniarz A. Molecular contamination of an animal facility during and after African swine fever virus infection. *J Vet Res.* 2023;67(4):503-8.
- Wang L, Kim J, Kang H, Park HJ, Lee MJ, Hong SH, Seo CW, Madera R, Li Y, Craig A, Retallick J, Matias-Ferreira F, Sohn EJ, Shi J. Development and evaluation of two rapid lateral flow assays for on-site detection of African swine fever virus. *Front Microbiol.* 2024;15:1429808.
- World Organization for Animal Health [WOAH]. Manual of diagnostic tests and vaccines for terrestrial animals [online]. Paris: WOAH; 2022. African swine fever. Available at: <https://www.woah.org/en/what-we-do/standards/codes-and-manuals/>. Accessed 5 May 2025.
- World Organization for Animal Health [OIE]. World Animal Health Information Database (WAHIS) Interface [database online]. African swine fever. Paris: OIE; 2019. Available at: [http://www.oie.int/wahis\\_2/public/wahid.php/Wahidhome/Home](http://www.oie.int/wahis_2/public/wahid.php/Wahidhome/Home). Accessed 10 June 2019.
- Xu Q, Li D, Chen X, Liu X, Cao H, Wang H, Wu H, Cheng T, Ren W, Xu F, He Q, Yu X, Li W. *In vivo* study of inoculation approaches and pathogenicity in African swine fever. *Vet Sci.* 2024;11(9):403.
- Yamada M, Masujin K, Kameyama KI, Yamazoe R, Kubo T, Iwata K, Tamura A, Hibi H, Shiratori T, Koizumi S, Ohashi K, Ikezawa M, Kokuho T, Yamakawa M. Experimental infection of pigs with different doses of the African swine fever virus Armenia 07 strain by intramuscular injection and direct contact. *J Vet Med Sci.* 2021;82(12):1835-45.
- Yoon H, Hong SK, Lee I, Choi DS, Lee JH, Lee E, Wee SH. Arthropods as potential vectors of African swine fever virus outbreaks in pig farms in the Republic of Korea. *Vet Med Sci.* 2021;7(5):1841-4.
- Zani L, Forth JH, Forth L, Nurmoja I, Leidenberger S, Henke J, Carlson J, Breidenstein C, Viltrop A, Höper D, Sauter-Louis C, Beer M, Blome S. Deletion at the 5'-end of Estonian ASFV strains associated with an attenuated phenotype. *Sci Rep.* 2018;8(1):6510.
- Zani L, Masiulis M, Bušauskas P, Dietze K, Pridotkas G, Globig A, Blome S, Mettenleiter T, Depner K, Karvelienė B. African swine fever virus survival in buried wild boar carcasses. *Transbound Emerg Dis.* 2020;67(5):2086-92.
- Zhai SL, Wei WK, Sun MF, Lv DH, Xu ZH. African swine fever spread in China. *Vet Rec.* 2019;184(18):559.
- Zhang X, Zhou L, Ge X, Gao P, Zhou Q, Han J, Guo X, Zhang Y, Yang H. Advances in the diagnostic techniques of African swine fever. *Virology.* 2025;603:110351.
- Zhao D, Sun E, Huang L, Ding L, Zhu Y, Zhang J, Shen D, Zhang X, Zhang Z, Ren T, Wang W, Li F, He X, Bu Z. Highly lethal genotype I and II recombinant African swine fever viruses detected in pigs. *Nat Commun.* 2023;14(1):3096.
- Zhou X, Li N, Luo Y, Liu Y, Miao F, Chen T, Zhang S, Cao P, Li X, Tian K, Qiu HJ, Hu R. Emergence of African swine fever in China, 2018. *Transbound Emerg Dis.* 2018;65(6):1482-4.
- Zsak L, Borca MV, Risatti GR, Zsak A, French RA, Lu Z, Kutish GF, Neilan JG, Callahan JD, Nelson WM, Rock DL. Preclinical diagnosis of African swine fever in contact-exposed swine by a real-time PCR assay. *J Clin Microbiol.* 2005;43:112-9.

\*Link is defunct