

Zoonotic Tuberculosis in Mammals, including Bovine and Caprine Tuberculosis

Infections caused by *Mycobacterium bovis*, *M. caprae*, *M. pinnipedii*, *M. orygis* and *M. microti*

Last Updated: October 2019



The Center for
Food Security
& Public Health



INSTITUTE FOR
INTERNATIONAL
COOPERATION IN
ANIMAL BIOLOGICS

IOWA STATE UNIVERSITY
College of Veterinary Medicine



OIE Collaborating Centre for
• Diagnosis of Animal Disease and
Vaccine Evaluation in the Americas
• Day-One Veterinary Competencies
and Continuing Education



Importance

Several closely related bacteria in the *Mycobacterium tuberculosis* complex cause tuberculosis in mammals. Each organism is adapted to one or more hosts, but can also cause disease in other species. The two agents usually found in domestic animals are *M. bovis*, which causes bovine tuberculosis, and *M. caprae*, which is adapted to goats but also circulates in some cattle herds. Both cause economic losses in livestock from deaths, disease, lost productivity and trade restrictions. They can also affect other animals including pets, zoo animals and free-living wildlife. *M. bovis* is reported to cause serious issues in some wildlife, such as lions (*Panthera leo*) in Africa or endangered Iberian lynx (*Lynx pardinus*). Three organisms that circulate in wildlife, *M. pinnipedii*, *M. orygis* and *M. microti*, are found occasionally in livestock, pets and people.

In the past, *M. bovis* was an important cause of tuberculosis in humans worldwide. It was especially common in children who drank unpasteurized milk. The advent of pasteurization, followed by the establishment of control programs in cattle, have made clinical cases uncommon in many countries. Nevertheless, this disease is still a concern: it remains an important zoonosis in some impoverished nations, while wildlife reservoirs can prevent complete eradication in developed countries. *M. caprae* has also emerged as an issue in some areas. This organism is now responsible for a significant percentage of the human tuberculosis cases in some European countries where *M. bovis* has been controlled. It was also found recently in China and North Africa, suggesting its distribution probably extends beyond Europe.

Etiology

In mammals, tuberculosis is caused by members of the *Mycobacterium tuberculosis* complex, which are Gram positive, acid-fast bacterial rods in the family Mycobacteriaceae. The organisms maintained in animals include *Mycobacterium bovis* (bovine tuberculosis), *M. caprae* (caprine tuberculosis), *M. pinnipedii*, *M. orygis* and *M. microti*. *M. caprae*, *M. pinnipedii* and *M. orygis* were members of *M. bovis* before being designated separate species. Two other agents, *M. tuberculosis* and *M. africanum*, are maintained in humans but occasionally affect animals. One poorly-understood organism, *M. canettii*, has only been found in humans in Africa, but it is not thought to be maintained in people. Its reservoir is unknown. Some authors also recognize additional species, such as *M. mungi* in banded mongooses (*Mungos mungo*) or *M. suricattae* in meerkats (*Suricata suricatta*).

The taxonomy of the *M. tuberculosis* complex can be controversial, and *M. bovis* and *M. caprae* are sometimes called *M. bovis* subsp. *bovis* and *M. bovis* subsp. *caprae*, respectively. Other authors argue that all of the agents in the *M. tuberculosis* complex should be considered a single species, due to their close genetic relationships. Under this system, *M. bovis* and *M. caprae* would be renamed *M. tuberculosis* subsp. *bovis* and *M. tuberculosis* subsp. *caprae*.

Species Affected

Mycobacterium bovis

Cattle are the primary hosts for *M. bovis*, but it can probably infect most or all mammals and marsupials. Susceptibility to illness might vary. Clinical cases have been described in sheep, goats and their relatives; pigs and other Suidae; horses and other equids; camels and South American camelids; dogs and other canids; cats and various wild felids; many wild ruminants and cervids; other ungulates such as elephants, rhinoceroses (*Ceratotherium simum*, *Diceros bicornis*) and giraffes (*Giraffa camelopardalis*); and diverse other hosts including ferrets, mink, banded mongooses, meerkats, otters (*Lutra lutra*), raccoons (*Procyon lotor*), coatis (*Nasua nasua*), opossums, hares (*Lepus* sp.), rabbits, nonhuman primates, bears and many rodents. Although most organisms previously reported in seals and sea lions are now classified as *M. pinnipedii*, *M. bovis* has been identified in grey seals (*Halichoerus grypus*), suggesting that marine mammals are probably susceptible to it as well.

Bovine Tuberculosis

Cattle are the usual maintenance hosts for *M. bovis*, but it can also be maintained in goats, captive cervids and some free-living wildlife. A species can be a maintenance host in one location and a spillover host at another, depending on its population density and other factors that facilitate transmission. Elk (*Cervus canadensis*) and white-tailed deer (*Odocoileus virginianus*) are both thought to be maintenance hosts around Riding Mountain National Park, Canada, though elk play the principal role. However, elk are spillover hosts in Michigan, where white-tailed deer maintain *M. bovis*. Neither species currently seems to be infected with this organism in other parts of the U.S. or Canada. Similarly, wild boar are thought to be important maintenance hosts in Mediterranean climates on the Iberian Peninsula in Europe, but they seem to be spillover hosts in Atlantic Spain and some other parts of Europe. Other maintenance hosts include wood bison (*Bison bison athabascae*) around Wood Bison National Park, Canada, feral swine in parts of Hawaii, brush-tailed opossums (*Trichosurus vulpecula*) in New Zealand, European badgers (*Meles meles*) in the United Kingdom and Republic of Ireland, red deer/elk (*Cervus elaphus*) in parts of Europe; and African buffalo (*Syncerus caffer*) and Kafua lechwe (*Kobus lechwe kafuensis*) in Africa. Greater kudu (*Tragelaphus strepsiceros*) and warthogs (*Phacochoerus aethiopicus*) have also been suggested as possible maintenance hosts in Africa, but their role is still debated. Determining whether a wildlife species is a maintenance host or a spillover host can be difficult. At one time, feral ferrets (*Mustela furo*) were proposed as maintenance hosts in New Zealand, but they now appear to be spillover hosts in which *M. bovis* can cycle for a long time but eventually disappears. Ecosystems where *M. bovis* circulates in wildlife usually have diverse spillover hosts ranging from rodents and small mammals to carnivores and ungulates.

Tuberculosis in birds is usually caused by *M. avium*, which is not a member of the *M. tuberculosis* complex. However, there are a few reports of birds affected by *M. bovis*, including parrots, an ostrich and a black swan (*Cygnus atratus*). *M. bovis* replicated in pigeons (*Columba livia*) after oral or intratracheal inoculation, and budgerigars (*Melopsittacus undulatus*) after intramuscular inoculation. Clinical signs developed in the budgerigars but not the pigeons; however, a few pigeons occasionally shed the organism in feces. American crows (*Corvus brachyrhynchos*), starlings (*Sturnus vulgaris*), Mallard ducks (*Anas platyrhynchos*) and wild turkeys (*Meleagris gallopavo*) seem to be resistant to experimental infection.

Mycobacterium caprae

M. caprae is an important cause of tuberculosis in goats. Clinical cases or infections have also been documented in cattle, sheep, pigs, wild boar, red deer, red foxes (*Vulpes vulpes*), and some animals in zoos including a Siberian tiger (*Panthera tigris altaica*), wild Bactrian camels (*Camelus bactrianus ferus*), a dromedary camel

(*Camelus dromedarius*), a Borneo elephant (*Elephas maximus borneensis*) and American bison (*Bison bison*). Although goats are the usual maintenance hosts, *M. caprae* has been found in cattle herds that have no apparent contact with small ruminants. In Europe, it seems to be maintained in free-living red deer and possibly in wild boar.

Mycobacterium microti*, *M. pinnipedii* and *M. orygis

M. microti is maintained in wild rodents and insectivores (e.g., shrews), and causes tuberculosis in these species. Commonly infected species such as field voles (*Microtus agrestis*), bank voles (*Myodes glareolus*), wood mice (*Apodemus sylvaticus*) and shrews (*Sorex araneus*) might be maintenance hosts. Wild boar are also proposed to maintain this organism. Some recognized spillover hosts include cattle, goats, pigs, South American camelids, cats, dogs and ferrets, as well as free-living or captive wildlife such as badgers, meerkats, otters and nonhuman primates (squirrel monkeys, *Saimiri sciureus*; marmosets, *Callithrix jacchus*).

M. pinnipedii, which causes tuberculosis in pinnipeds, has been isolated from several species of seals and sea lions. There are occasional reports of this organism in other hosts including cetaceans (Hector's dolphin, *Cephalorhynchus hectori*; bottlenose dolphin, *Tursiops truncatus*), cattle, llamas, Bactrian camels, at least two species of tapirs (*Tapirus* sp.) and nonhuman primates (e.g., gorillas, *Gorilla* sp.). Mycobacterial infections in a captive snow leopard (*Panthera uncia*) and Amur leopard (*Panthera pardus orientalis*) at one zoo may also have been caused by *M. pinnipedii*.

M. orygis was first isolated from East African oryx (*Oryx beisa*). Since then, it has been found in a number of other ungulates including cattle, Arabian oryx (*Oryx leucoryx*), waterbuck (*Kobus ellipsiprymnus*), African buffalo, Indian rhinoceros (*Rhinoceros unicornis*), spotted deer (*Axis axis*), blue bull (*Boselaphus tragocamelus*) and nonhuman primates.

Zoonotic potential

M. bovis, *M. caprae*, *M. orygis*, *M. pinnipedii* and *M. microti* can all cause tuberculosis in humans.

Geographic Distribution

A limited number of countries (e.g., Australia, Iceland, Greenland, Singapore, some European nations, Israel) are completely free of *M. bovis*; however, infected livestock herds are now uncommon in Europe, Canada, the United States, New Zealand and some other locations. Wildlife reservoirs are known to exist in Wood Bison National Park and Riding Mountain National Park in Canada, Hawaii and northeastern Michigan in the U.S., some European countries and New Zealand. Bovine tuberculosis is still common or relatively common in cattle in parts of Africa, Asia, the Middle East and Latin America including Mexico.

Bovine Tuberculosis

M. caprae has mainly been reported in Europe; however, it was recently described in a few cattle in North Africa (Algeria, Tunisia), and sheep and reindeer in China, suggesting that it may be more widely distributed. *M. orygis* has been found in Africa, southern Asia and the Middle East. Isolated cases identified in humans in the U.S. and New Zealand were probably acquired in Asia. *M. pinnipedii* has been described in marine mammals in many locations, including the coasts of Europe, New Zealand, Australia and South America. *M. microti* also seems to be widespread, with reports of indigenously acquired clinical cases in Europe, Africa and South America. It seems likely that some of these organisms also circulate in other areas.

Transmission

Most information on the transmission of the zoonotic mycobacteria comes from studies of *M. bovis*. Depending on the sites where it has localized, this organism may be found in respiratory secretions, exudates from lesions (e.g., draining lymph nodes, some skin lesions), urine, feces, milk, vaginal secretions and semen. Shedding can occur intermittently and differs between individuals. *M. bovis* is more likely to be transmitted when the respiratory tract is affected and in the late stages of the disease, when lesions are more extensive. Close contact in confined spaces facilitates its spread. It has also been cultured from the oral secretions of some animals, including ferrets, which may facilitate transmission in bites.

Animals can become infected by inhalation, ingestion or direct contact through the mucous membranes or breaks in the skin. Much larger numbers of organisms are generally needed to establish an infection by ingestion than inhalation. The importance of the various transmission routes differs between host species. Cattle are often infected via aerosols during close contact. Ingestion is less important in this species, except in calves that nurse from infected cows. Cutaneous, genital (sexual) and congenital transmission are possible but seem to be uncommon in cattle. Respiratory transmission is also thought to predominate in some other hosts such as camels, nonhuman primates and badgers. However, ingestion is thought to be the most common route in ferrets, cats, deer and horses. Percutaneous transmission is mainly seen in species that tend to hunt or fight, such as cats and badgers. Cats might also inoculate organisms onto mucous membranes when they wash after eating. Dogs with kidney lesions were thought to have transmitted *M. bovis* in urine in a kennel. Nosocomial transmission has been reported in at least two small animal veterinary clinics. In one instance, organisms from an infected cat were transmitted to healthy feline surgical patients, probably on hands or clothing.

Humans can be infected by the same routes as animals, including inhalation, ingestion and direct contact with mucous membranes or breaks in the skin. Ingestion is a common form of exposure when dairy products are not pasteurized. Viable *M. bovis* may also be found in raw or undercooked meat and other animal tissues. Animal bites have been linked rarely to human cases. Person-to-person

transmission is possible, especially when the respiratory tract is affected. These incidents usually involve people in close contact, such as families, but at least one investigation suggested the possibility of respiratory transmission of *M. bovis* during casual contact in the community. Possible human-to-human transmission was also suggested for *M. microti*. Rare cases of probable or possible human-to-animal transmission have been documented for *M. bovis* and *M. orygis*. The organism was probably transmitted in aerosols in most of these cases, but urine-contaminated hay was thought to be involved in one incident in cattle.

The environmental survival of *M. bovis* and closely related mycobacteria are affected by the temperature, humidity, exposure to sunlight, fluctuations in moisture and temperature, competition with other microorganisms and initial numbers of bacteria. *M. bovis* can persist for up to several months in soil and other material (e.g., feed, feces), particularly in cold, dark and moist conditions; however, it sometimes disappears within days or weeks, especially when exposed to direct sunlight, higher temperatures and dry environments. There are a few reports of extended survival for a year or more, generally in feces or soil under optimal laboratory conditions. One study, which tested soil, water, hay and shelled corn kept outdoors in open containers, found that the number of viable *M. bovis* declined significantly during the first 2 weeks, but a small number of organisms survived longer. *M. bovis* is infrequently isolated from soil or pastures grazed by infected cattle, but whether this is due to inactivation of the organism or difficulties in isolating it from sites with competing bacteria, which grow more readily in culture, is not always clear.

Disinfection

The members of the *M. tuberculosis* complex are relatively resistant to disinfectants. Some agents effective against *M. tuberculosis*, which is more resistant than *M. bovis*, include phenol-based disinfectants, povidone iodine (but not iodophors), sodium hypochlorite, peracetic acid, glutaraldehyde, formaldehyde, orthophthaldehyde ethylene oxide, and a mixture of 7.5% hydrogen peroxide and 0.85% phosphoric acid. Some of these agents may require a longer contact time or higher concentrations than normally used for other microorganisms. In one report, a 70% alcohol solution could only inactivate *M. tuberculosis* in the absence of sputum; however, alcohols may be a valuable component in multi-step disinfection procedures for equipment.

M. bovis and *M. caprae* are both destroyed by standard HTST pasteurization of milk. Mycobacteria can also be killed by exposure to UV light, heating to greater than 65°C (149°F) for at least 30 minutes, or autoclaving (moist heat of 121°C/ 250°F for at least 15 minutes).

Infections in Animals

Incubation Period

Tuberculosis usually has a slow onset, with clinical signs often taking several months or more to develop. Infections can also remain latent for years and later reactivate. Unusually short incubation periods have been seen occasionally, including in 2 cats accidentally infected via surgical incisions, which became ill after 14 and 42 days.

Clinical Signs

Bovine tuberculosis (Mycobacterium bovis)

Tuberculosis is usually a chronic, debilitating disease in cattle, although acute and rapidly progressive cases are possible. The onset is typically insidious, initially with few or no signs of illness. Common clinical signs in this species include weight loss (emaciation may become severe in the terminal stages), weakness, inappetence, a low-grade, fluctuating fever, lymphadenopathy and respiratory involvement with a moist, intermittent cough that is worse in the morning, or during cold weather or exercise. Dyspnea or tachypnea may also be seen, and involvement of the gastrointestinal tract can result in intermittent diarrhea and constipation. The superficial lymph nodes can be palpably enlarged, and sometimes rupture and drain. The retropharyngeal lymph node is often affected in this species. Enlargement of deeper lymph nodes occasionally obstructs blood vessels, airways or the digestive tract. Ocular disease (e.g., anterior uveitis, choroidal infiltrates, subretinal exudates), repeated abortions and infertility caused by lesions in the uterus, and other syndromes have also been reported. Skin lesions are uncommon in cattle.

Tuberculosis is generally similar in other species, but the predominant syndromes or course of the disease may differ. Abdominal organs are affected more often than the respiratory tract in equids and some other hosts, while elephants tend to have few signs until the lesions are extensive. Syndromes reported in farmed cervids range from cases that progress slowly, over a period of years, with abscesses of unknown origin in isolated lymph nodes, to disseminated disease with a rapid, fulminating course. Skin lesions are sometimes seen in cervids. Subclinical infections seem to be prevalent in pigs, but disseminated disease can also be seen, especially in young animals. Osteomyelitis and meningeal involvement are reported to be relatively common in this species. Respiratory disease is a common form of tuberculosis in both badgers and brush-tailed opossums, but it tends to be chronic in badgers, which may survive for years, while opossums often die within a few months. Wildlife with advanced tuberculosis are sometimes found when they display abnormal behavior (e.g., opossums wandering around in the daytime).

Localized or generalized tuberculosis can be seen in cats. The submandibular lymph node is often enlarged in this species. In addition to lymphadenopathy, cats with

generalized disease usually have nonspecific signs of illness (e.g., weight loss, persistent or fluctuating low-grade fever) and, in some cases, respiratory signs (coughing, dyspnea) or episodes of vomiting or diarrhea. Central nervous system (CNS) signs, enlargement of abdominal organs (e.g., liver, spleen) and joint or bone involvement are also possible. Skin and subcutaneous lesions are common in cats, and may appear as a soft swelling, firm nodule or flat ulcer. Skin lesions often result from direct inoculation of the organism, and tend to affect the face, neck, shoulders and extremities. Progressive lesions may sometimes expose and destroy the bones of the nose and face. Some infected cats have unilateral or bilateral eye lesions, either alone or accompanied by other syndromes. Common ocular lesions include panuveitis or anterior uveitis, conjunctivitis, and granulomas in ocular tissues (e.g., cornea, choroid, retina), the eyelid or periorbital tissues. As in other species, tuberculosis normally progresses slowly in cats. However, fulminant disease is possible. In one instance, two cats accidentally infected through their surgical wounds became severely ill within 2-3 weeks. Tuberculosis in lions seems to resemble the illness in cats.

There are usually no distinctive features of bovine tuberculosis in dogs, but one outbreak in a foxhound kennel was characterized by an unusual fulminant course with nonspecific signs including weight loss and lymphadenopathy, together with polyuria and polydipsia from kidney involvement.

Clinical cases caused by *M. bovis* are rare in birds, and few descriptions are available. Sudden death was reported in a black swan affected by this organism, and nonspecific signs of illness were seen in experimentally infected birds. This appears similar to avian tuberculosis, which is caused by organisms such as *M. avium* and is characterized mainly by poor body condition and other nonspecific signs, although skin lesions, ocular involvement and signs related to lesions in the internal organs are also possible.

Tuberculosis caused by other organisms

In terrestrial mammals, clinical cases caused by other organisms are usually indistinguishable from those caused by *M. bovis*. However, a clinical picture that consists of firm, raised skin lesions and submandibular lymphadenopathy is common in cats infected with *M. microti*, either with or without the involvement of other organs. *M. microti* can also cause other syndromes in cats, including pulmonary disease, arthritis, ocular signs and disseminated disease. Lethargy, anorexia, weight loss and respiratory signs are common in pinnipeds infected with *M. pinnipedii*.

Post Mortem Lesions [Click to view images](#)

Mammals

Tuberculosis is characterized by the formation of granulomas where bacteria have localized. The classic lesion is the tubercle, a yellowish-white or grayish-white

granuloma typically enclosed in a capsule of varying thickness. Its interior is usually caseous, caseo-calcareous or calcified in cattle, small ruminants, pigs and a number of other species. However, tubercles in some species of cervids are often poorly encapsulated, tend to resemble abscesses and may have purulent centers. Tubercles vary in size. Some are small enough to be missed unless the tissue is sectioned; others may form coalescing, confluent lesions that affect most of the organ. In cattle and other species that tend to inhale the organisms, lesions are typically found in the lungs and lymph nodes of the head and thorax. They are more likely to occur in the abdominal organs and lymph nodes (e.g., mesenteric lymph nodes) of animals infected by ingestion. Many other tissues and organs including the skin, bones, joints and CNS can also be affected.

Classic tubercles are less common in some hosts such as cats, dogs and horses. Instead, the lesions tend to appear as diffuse granulomatous infiltration, which can resemble neoplasia, or multiple small uncalcified nodules or caseous foci. Smooth, gray infiltrative lesions in the spleen and liver, are characteristic of primary tuberculosis lesions in horses. Small caseous foci may sometimes be found within these lesions. Diffuse granulomatous infiltrates are also common in disseminated disease, including cases in hosts that normally form classic tubercles at the primary infection site. Miliary tuberculosis, the most common form of disseminated disease in cattle, appears as small gray, white or yellowish-white caseous foci scattered throughout a tissue.

In cattle, dissemination of *M. bovis* to the pleura, pericardium or peritoneum results in multiple small, sessile, pedunculated or cauliflower-like tubercles, which eventually calcify. They are said to resemble pearls, and can sometimes be so numerous that they cover the serosa. Dissemination to serosal surfaces is reported to be infrequent in sheep, goats and pigs, but horses sometimes have nodular serosal lesions. Pleuritis, peritonitis and pericarditis also be seen in some cats and dogs, but the serosa is usually thickened by granulation tissue rather than forming nodules. Some other lesions that may be found in tuberculosis include straw-colored or serohemorrhagic fluid in the body cavities or pericardium; ulcerative lesions in the trachea, bronchi or intestines; osteomyelitis; and various abnormalities in the joints, including proliferative lesions and joint capsule mineralization. Some infected animals do not have any visible lesions.

Birds

Mycobacterium sp. do not form classic encapsulated tubercles in birds; instead, the internal organs usually contain pale yellow, white or tan nodules and granulomatous infiltration. Small nodules are usually not caseated; larger nodules may be caseous but do not usually mineralize. Some birds with mycobacteriosis have no visible lesions.

Diagnostic Tests

Tuberculosis can be diagnosed by detecting either the causative organisms or the immune responses to these organisms. Animals that react in either type of test are generally treated as having an active infection. In some species, diagnostic tests may be supplemented with x-rays or other imaging techniques to visualize abnormalities in the internal organs.

Tuberculin test

The tuberculin skin test, which detects cell-mediated immune responses (CMI), is the primary screening test in cattle and some other species. Tuberculin, a mixture of bacterial proteins, is injected intradermally and the site is examined 48-96 hours later for an inflammatory swelling (delayed hypersensitivity reaction). The elapsed time affects the test's sensitivity and specificity. Livestock tests normally use antigens from *M. bovis*, but they can also detect infections with other members of the *M. tuberculosis* complex. A comparative test that evaluates relative reactivity to *M. bovis* and environmental mycobacteria (*M. avium*) can be used to decrease false positive reactions. False negative tests are also possible, especially during the first few weeks after exposure, in animals with suppressed immune responses, including those that have recently calved, and late in the course of the disease, when CMI may diminish.

The primary tuberculin tests used in cattle are the caudal fold test (CFT), the single cervical intradermal test (CIT) and the comparative cervical test (CCT). The CIT is generally considered to have higher sensitivity and specificity than the CFT; however, the CFT uses a site that can be more convenient to inject and examine. The U.S., Canada and some other countries use the CFT for preliminary screening of cattle, and re-test reactors with the CCT. Some European countries use the CCT; others start with the CIT and confirm reactors with the CCT.

The tuberculin test is also used in other animals, but it must be validated in each species, and the injection site can differ. The CCT and CIT are common tests for tuberculosis in captive cervids. Tuberculin tests have also been evaluated and/ or used in goats, water buffalo, guinea pigs and a limited number of wildlife hosts (e.g., lions infected with *M. bovis*, captive sea lions infected with *M. pinnipedii*). There is relatively little research on these tests in sheep, and some authors report that false negatives are common unless the animal has extensive lesions. Tuberculin tests have relatively poor sensitivity and specificity in llamas and alpacas, and they are often supplemented with *in vitro* CMI tests or serology for better accuracy. They are considered to be unreliable, due to inadequate sensitivity and/or specificity, in cats, dogs, horses, European badgers and brush-tailed opossums. Practical considerations, including the need to recapture the animal to read the test, can also preclude their use.

Tuberculin tests cannot be employed in pachyderms due to the nature of their skin.

In vitro tests of cell-mediated immunity

Gamma-interferon (IFN- γ) release assays measure CMI using blood samples. Most of these tests are in the ELISA or ELISPOT format. Comparative assays that evaluate reactivity to *M. bovis* and environmental mycobacteria (*M. avium*) may be available in some laboratories. IFN- γ release tests can be used instead of skin tests in animals that are difficult to capture or handle. They have also been combined with skin tests to increase sensitivity or specificity.

IFN- γ release tests developed for cattle do not necessarily work in other animals. As of 2019, these tests have been used or evaluated in a limited number of species including goats, sheep, South American camelids, cats, badgers and lions. They were also employed, in conjunction with serology, during an outbreak in dogs. Samples for this test must be transported to the laboratory promptly, as it must be started within 24-30 hours of blood collection.

The lymphocyte proliferation test, another blood test for CMI, has mainly been used in wildlife and zoo animals. It is not a commonly used test.

Serology

Antibodies to *M. bovis* usually develop later than CMI, and titers tend to increase as the disease progresses. The most commonly used serological tests are ELISAs and lateral flow immunochromatographic assays, but other methods (e.g., immunoblotting or a multiantigen print assay) may also be available. Serological tests have only been validated for a small number of hosts. However, they have been used to help diagnose tuberculosis in South American camelids, farmed cervids, dogs, elephants, and a number of captive or free-living wildlife infected with *M. bovis*, and in pinnipeds infected with *M. pinnipedii*. At one zoo, gradually increasing antibody titers preceded positive tuberculin skin tests in a *M. bovis*-infected onager (*Equus hemionus onager*) and gemsbok (*Oryx gazelle gazelle*). Serology may occasionally be useful in cattle or goats in the late stages of tuberculosis. There are a few reports of these tests being used to monitor responses to treatment (e.g., in elephants). Cross reactions to other bacteria, especially environmental mycobacteria, can complicate test interpretation.

Tuberculin skin tests can boost serological responses in deer and South American camelids, and combining these tests can reveal infected animals that do not respond in the skin test. This technique needs to be validated in each species. An attempt to use it in elephants resulted in a high rate of false positive reactions.

Detection of the organism

In live animals, members of the *M. tuberculosis* complex are sometimes found in exudates, biopsy samples from affected tissues, sputum and other secretions and excretions. Trunk washes to collect sputum are a routine diagnostic test in elephants, as there are few other

antemortem tests in this species. Sputum is also collected in captive pinnipeds that may be infected with *M. pinnipedii*, and bronchoalveolar lavage fluid has occasionally been employed in lions or other animals. More often, tuberculosis is confirmed in tissue samples from affected organs, taken at necropsy. Infections can sometimes be found in animals with no apparent lesions. In cattle with no visible lesions, the recommended minimum sample is pooled lymph nodes from the head and thorax. Collecting additional tissues may be helpful. A recent study found that, in cattle with no visible lesions that were infected by ingestion, *M. bovis* is more likely to be found in the abdominal (e.g., mesenteric) lymph nodes.

A presumptive diagnosis can be made by histopathology and/or the demonstration of acid-fast bacilli in smears from tissues, exudates or body fluids. Some lesions contain abundant bacteria; others have very few. Concentration methods can increase the chance of visualizing the organism. Direct smears for microscopy can be stained with Ziehl-Neelsen stain, a fluorescent acid-fast stain (e.g., auramine) or immunoperoxidase techniques. *M. microti* is distinctive in that the rods may appear curved in fresh specimens. This characteristic is usually lost after culture.

Tuberculosis can be confirmed by isolating the causative organism on selective culture media such as modified Middlebrook 7H10 or 7H11 agar, or Stonebrink's or Lowenstein-Jensen egg-based medium. While colonies may occasionally appear as early as 2 weeks, the median time on solid media is 4-5 weeks, and some samples may require up to 12 weeks or longer. Recovery is reported to be faster in automated liquid broth culture systems. Because these slow-growing organisms can be overgrown by contaminants, samples for culture should be collected as aseptically as possible. If necessary, culture can incorporate a decontamination step, where the samples are treated with a toxic agent to which members of the *M. tuberculosis* complex are relatively resistant (e.g., cetylpyridinium chloride, sodium hydroxide or oxalic acid). However, decontamination also decreases the recovery of mycobacteria. Culture is not always successful in infected animals. *M. microti* is especially difficult to isolate.

A cultured organism's identity can be confirmed with biochemical tests, or PCR and other genetic techniques (e.g., commercial DNA probes; sequencing of certain genes). Biochemical methods are slow and labor-intensive and may occasionally give ambiguous results, and molecular methods are generally preferred if they are available. At least one commercial genetic test can distinguish *M. bovis*, the vaccine strain *M. bovis* BCG, *M. caprae*, *M. tuberculosis*, *M. africanum* and *M. microti*. However, it cannot reliably differentiate some organisms, such as *M. pinnipedii* and *M. africanum*. Genetic techniques such as spoligotyping, multilocus variable number tandem repeat typing (MLVA) or whole genome sequencing can be used for epidemiological purposes such as tracing outbreaks.

PCR can sometimes identify the causative organism directly in clinical samples. It has the advantage of speed, and can also be helpful when organisms are difficult to grow. However, it is less sensitive than culture if the sample has few bacteria.

Animal inoculation is possible but rarely done.

Treatment

Antibiotics have been used to treat some animals with tuberculosis, especially pets and zoo animals. However, the possibility of clinical improvement without bacteriological cure must be kept in mind. Some animals that responded initially later relapsed, especially with inadequate treatment (e.g., too short treatment or the use of a single drug). The risk of shedding organisms, hazards to humans (especially if the respiratory tract is infected or there are draining lesions), and potential for the development of drug resistance make treatment controversial. Treatment is not permitted in some countries.

Members of the *M. tuberculosis* complex are not susceptible to many common antibiotics; they can be treated only with a limited number of tuberculocidal agents. Treatments for animals are usually patterned after successful protocols used in humans, and employ two or more drugs, given simultaneously for months. When selecting antibiotics, it should be kept in mind that, except in rare cases, *M. bovis* is intrinsically resistant to pyrazinamide, a common “first line” drug for tuberculosis. This drug can, however, be used in animals infected with *M. caprae*. One protocol for cats, which are unusually sensitive to some of the commonly used tuberculocidal drugs, combines rifampicin, a fourth generation fluoroquinolone such as pradofloxacin, and clarithromycin or azithromycin.

Surgery is occasionally used to remove small masses, treat ocular tuberculosis (e.g., enucleation), or amputate an infected joint, in combination with tuberculocidal drugs. Drug treatment alone seems to have poor results in cats that have significant osteomyelitis or joint involvement.

Control

Disease reporting

Veterinarians who suspect an animal is infected with a member of the *M. tuberculosis* complex should follow their national and/or local guidelines for disease reporting. Although most regulations are written for *M. bovis* or *M. caprae* in livestock, some nations stipulate that other members of this complex or infected pets (e.g., cats with tuberculosis in the U.K.) must also be reported. State authorities should be consulted for regulations in the U.S.

Prevention

Sanitation and disinfection, open air housing rather than confinement, and avoidance of crowding might reduce the spread of tuberculosis within a herd. Rodent control decreases the risk that these animals may disseminate the

organism. Keeping cats indoors can help protect them in areas endemic for *M. bovis*, *M. caprae* or *M. microti*. In endemic areas, barrier nursing precautions and tuberculocidal disinfectants should be considered in veterinary hospitals when animals have signs consistent with tuberculosis.

Control programs are usually targeted at bovine tuberculosis in cattle, but some programs may include *M. caprae*. In these programs, animals are tested periodically with the tuberculin skin test and/or other assays. If an infected herd is found, the reactors are removed and the herd is quarantined until all animals test negative. Where testing is done regularly, most herds contain only a few infected cattle. Reactors are generally slaughtered, but some countries may employ test-and-segregation programs at first, and later switch to test-and-slaughter. Entire herds can also be depopulated, though this is generally rare. Once eradication is nearly complete, slaughter surveillance, with tracing of infected animals, may be a more efficient use of resources. However, it is relatively insensitive, and can miss infected animals with few or no visible lesions. Screening programs may sometimes include species other than cattle (e.g., farmed deer); however, in most cases, these animals are only found by passive surveillance or epidemiological investigations of infected cattle.

The occurrence of *M. bovis* and *M. caprae* in wildlife reservoirs complicates eradication efforts. Transmission from these animals to livestock can be reduced with biosecurity measures such as wildlife barriers around feed storage areas, or solid metal barriers and gates to exclude badgers from cattle pens. Some countries have also established control programs targeted at the principal maintenance host(s). Culling may decrease their population density below the level needed to sustain transmission. However, each situation must be assessed individually, as culling may have unanticipated effects if it encourages infected animals to disperse. Capture and testing programs, with the release of uninfected animals, have sometimes been used in badgers and African buffalo. Some control programs include bans on feeding deer and elk, to reduce transmission between congregating animals and discourage transmission on fomites. Elimination of organisms mainly found in wildlife, such as *M. pinnipedii*, *M. orygis* and *M. microti*, is impractical.

Vaccines, including some that contain the Bacillus Calmette Guerin (BCG) strain of *M. bovis*, are being evaluated for possible use in livestock and farmed deer, and as an aid in controlling tuberculosis in wildlife reservoirs. IFN- γ release assays and tuberculin skin tests that can distinguish BCG-vaccinated livestock from animals infected with *M. bovis* have been published. As of 2019, the only licensed vaccine is a parenteral BCG vaccine for badgers in the U.K. Test-negative badgers are vaccinated before release in some control programs.

Morbidity and Mortality

Possible outcomes after exposure to a member of the *M. tuberculosis* complex include elimination of the organism by the immune system, persistent asymptomatic infection with or without limited lesions, or the development of a chronic and eventually fatal disease. Some subclinically infected animals become ill only if they become debilitated or stressed, or in old age. There are also reports of *M. bovis* lesions resolving in experimentally infected animals, with or without the elimination of the organism; however, this is thought to be rare. Factors that may influence the outcome of an infection include the animal's species, the species of *Mycobacterium*, the dose of organisms, the site of inoculation and the individual's general health. Clinical cases also seem to be more severe in young animals. Genetic resistance to *M. bovis* has been identified in some hosts including cattle, farmed red deer and African buffalo. Zebu cattle (*Bos indicus*) are reported to be more resistant to bovine tuberculosis than the Ankole or Holstein breeds of *Bos taurus*.

Mycobacterium bovis

In two studies of *M. bovis* transmission in naturally infected cattle, 0-40% of susceptible contacts reacted in tuberculin tests, and 0-10% developed gross lesions. Sheep are thought to be more resistant to *M. bovis* than cattle or goats, but underdiagnosis could be an issue, as reports of tuberculosis in sheep seem to be rising. It is possible that management (e.g., less intensive rearing) or behavior helps protect sheep from exposure. Rabbits also seem to be relatively resistant; most clinical cases have been reported in animals exposed to heavily contaminated environments.

Tuberculosis was seen regularly in pigs, horses, cats and dogs in the first part of the 20th century, when infected cattle with extensive lesions were more common. Only sporadic cases have been reported in dogs or horses in recent years. During one recent outbreak in a working English Foxhound kennel, approximately 8% of the dogs were infected, and 8 of the 14 infected dogs became ill. Ferrets, guinea pigs and cats seem to be relatively susceptible to *M. bovis*. At one time, infections were estimated to occur in 7% of the cats in Switzerland, and about half of the cats exposed to infected cattle herds in Pennsylvania. Later, in the 1970s, a Swiss study found evidence of tuberculosis in 0.2% of feline necropsies. However, recent reports did not detect any infected cats on farms with bovine tuberculosis in Michigan or the U.K., probably because the number of infected cattle was low and the lesions were not extensive. In the U.K., infected rodents and other wildlife now seem to account for most cases of tuberculosis in cats. Occasional infections in indoor cats probably result from exposure to other infected cats or people in the household. A commercial raw food with venison was also implicated in a number of recent cases. Most cats with tuberculosis are not infected with immunosuppressive viruses, and were previously healthy.

M. bovis infection rates in wildlife spillover or maintenance hosts generally range from $\leq 5\%$ to 20%, but there are reports of localized clusters with higher prevalence. High animal density, social structures that encourage contact between individuals, and artificial concentration of animals (e.g., supplemental feeding of deer) increase the probability that a species will maintain *M. bovis*. Spillover hosts sometimes act as temporary reservoirs that can reinfect maintenance hosts.

Other organisms

Research on *M. caprae* is limited, but many aspects of its morbidity and mortality are probably similar to *M. bovis*. Some evidence suggests that *M. caprae* tends to be less virulent than *M. bovis* in goats, although it can also cause severe disease. *M. pinnipedii* is thought to be widespread in pinnipeds, and *M. microti* is common in some wild rodent populations. Among domestic animals, clinical cases and lesions caused by *M. microti* have been reported relatively often in cats and wild boar, and sporadically or rarely in some mammals such as cattle. This organism is challenging to culture and it may be underdiagnosed. Some reports suggest that tuberculosis in cats tends to be less severe when it is caused by *M. microti* than *M. bovis*.

Infections in Humans

Incubation Period

Systemic signs can develop months to years after exposure in humans, or the infection may remain latent until many years later when waning immunity allows the organisms to reactivate. Tuberculous chancre, a type of skin lesion, typically appears 2-4 weeks after cutaneous exposure.

Clinical Signs

In humans, tuberculosis is characterized by nonspecific signs such as fever, malaise and weight loss, and various symptoms caused by lesions in the lymph nodes, lungs, skin, bones, joints, genitourinary system, intestinal tract, CNS and/or a variety of less frequently involved sites. In one unusual case caused by *M. bovis*, the primary sign of infection was a tooth abscess. Pulmonary involvement, with fever, chronic weight loss, coughing, chest pain and hemoptysis, is common in people infected with the anthroponotic organism *M. tuberculosis*, which is usually acquired by inhalation. Because zoonotic organisms are often ingested, many of these cases tend to involve extrapulmonary sites instead. Cervical lymphadenopathy of the jugulodigastric (tonsillar) and preauricular lymph nodes is a common presentation in children who drink infected milk. These nodes sometimes suppurate and drain to the skin, resulting in chronic skin lesions. Pulmonary involvement is, however, common when zoonotic mycobacteria cause disseminated disease or the organisms in latent infections reactivate. It can also be seen when zoonotic organisms are inhaled.

Bovine Tuberculosis

The skin lesions that occur in some cases of tuberculosis have been given various names (e.g., scrofuloderma, lupus vulgaris) depending on their appearance, origin and typical course. They can result from direct inoculation of organisms into the skin, or by the extension of an infection from another site. Some are also caused by hypersensitivity reactions to small numbers of organisms in cutaneous blood vessels. Cutaneous lesions may appear as papules, ulcers and/or suppurative lesions; soft, gelatinous plaques with central atrophy; reddish-brown, gradually enlarging subcutaneous nodules; or vegetative lesions that can resemble a tumor. One common localized form of tuberculosis, called tuberculosis verrucosa cutis or “butcher’s wart,” results from direct inoculation of the organism into the skin, often on the extremities, and typically appears as one or more painless verrucous and tuberous papules. Tuberculosis verrucosa cutis usually evolves slowly, does not ordinarily spread to the lymph nodes or other organs, and tends to be self-limiting. Another localized form called tuberculous chancre, which appears as a shallow, painless, nonhealing ulcer, may regress spontaneously, persist or evolve into other forms, and sometimes spreads to other parts of the body.

CNS disease and disseminated disease are particularly serious forms of tuberculosis. Patients with CNS disease can have meningitis, meningoencephalitis or a focal infection, either with or without the involvement of other organs. Chronic meningitis, the most common form of CNS disease, has an insidious onset and is most often seen in young children, people who are immunosuppressed and older adults. Meningoencephalitis has been reported in all ages, and patients can deteriorate fairly quickly. Disseminated disease usually affects both pulmonary and extrapulmonary organs, and occasionally includes widespread skin lesions. Patients with this form of tuberculosis are severely ill, and may die soon after the onset of clinical signs. Disseminated tuberculosis is particularly common in immunocompromised patients, and also has an elevated incidence in young children.

Diagnostic Tests

Tuberculin skin tests, IFN- γ release assays, direct microscopy for acid-fast bacilli, culture, and PCR or other nucleic acid assays, combined with imaging studies such as chest x-rays, are used to diagnose tuberculosis in humans. In people, the tests that evaluate immune responses are interpreted in conjunction with evidence of bacterial replication and disease. Unlike in animals, the diagnosis is not a binary assessment of whether the person is infected or not, but also considers possibilities such as a latent infection or exposure without persistence of the bacterium. Vaccination with BCG or sensitization by periodic skin testing can complicate the interpretation of some tests of CMI. Comparative tests that evaluate immune responses to environmental mycobacteria are employed only rarely in humans.

Human diagnostic laboratories sometimes use PCR-based systems to screen sputum samples for tuberculosis, either routinely or when they contain acid-fast bacilli. These tests are more sensitive than microscopic examination of the sample, but less sensitive than culture. Zoonotic mycobacteria, which often cause extrapulmonary tuberculosis, are not always found in sputum. Diagnostic testing for tuberculosis in people does not necessarily identify the species of organism except as a member of the *M. tuberculosis* complex. This is especially common in some impoverished countries, where acid-fast smears are widely used for diagnosis, and culture may be unavailable. The inability to identify *M. bovis* might influence the effectiveness of treatment.

Treatment

Commonly-used first line agents for tuberculosis in humans include rifampicin, isoniazid and pyrazinamide; other (second-line) drugs are employed for drug-resistant isolates. Multidrug-resistant strains of *M. bovis* (defined as resistance to at least isoniazid and rifampicin) have been identified occasionally, and there have been a few strains resistant to second line drugs. Antibiotic resistant strains of *M. bovis* are mainly thought to arise during tuberculosis treatment in humans, rather than circulating in animals.

Active tuberculosis is usually treated for a minimum of 6 months with two or more agents, used simultaneously. The standard first line tuberculosis treatment for *M. tuberculosis* can usually be employed for cases caused by *M. caprae*; however, *M. bovis* is intrinsically resistant to pyrazinamide and the recommended length of treatment for this organism is currently 9 months. Compliance can be an issue, and some researchers have been investigating whether shorter treatment periods would also be effective. Surgery might occasionally be considered for some isolated forms of tuberculosis such as cutaneous disease.

Prevention

The risk of contracting zoonotic tuberculosis can be decreased by controlling the organisms in domestic animals. Pasteurization destroys the members of the *M. tuberculosis* complex in dairy products. Thorough cooking is effective in meat and other animal tissues; however, some organisms may survive in undercooked (e.g., rare) meat. Carcasses with lesions should not be eaten. People who must handle infected animals or their tissues should use gloves and other personal protective equipment (PPE), including respiratory protection where aerosolization could be an issue. Any open wounds should be covered. Hunters should also use PPE in areas where wildlife may be infected. The use of high pressure hoses for cleaning animal facilities has been associated with some infections, probably by aerosolizing bacteria.

Some countries administer a BCG vaccine, which contains an attenuated strain of *M. bovis*, to infants. It can

Bovine Tuberculosis

protect young children from severe tuberculosis, especially disseminated disease and CNS disease; however, its efficacy differs between individuals, the protection wanes early in adolescence, and revaccination does not seem to be helpful. The BCG vaccine is expected to be protective against *M. bovis* as well as *M. tuberculosis*.

Morbidity and Mortality

M. bovis

Current understanding of human tuberculosis suggests that, in some cases, the organisms are eliminated from the body after a few weeks, while, in others, they may proliferate either with or without clinical signs, or establish a latent infection. Infected individuals can move between these states. The typical course of events is for inapparent infections to become symptomatic, or latent infections to become reactivated when immunity wanes. However, spontaneous improvement or recovery is also possible, even after the development of clinical signs. While there are no data for *M. bovis*, approximately 5-10% of humans infected with *M. tuberculosis* become ill during the first 2 years. The risk of reactivation in latent *M. tuberculosis* infections is estimated to be 5-10% in healthy people over their lifetime, and about 10% per year if the person is significantly immunocompromised.

In healthy people, clinical tuberculosis is often a slowly progressive disease with low mortality if it is treated. However, disseminated disease and CNS involvement are more likely to be fatal. In a small number of studies, case fatality rates were 18-72% in various forms of CNS tuberculosis, and about a third of the surviving patients had permanent neurological deficits. Mortality rates are also elevated in people who are immunocompromised, regardless of treatment, and in young children. Childhood mortality from tuberculosis was about 1% in recent years. Without treatment, however, historical reports suggest case fatality rates of 22% in all children and 44% in those under the age of 5 years. Only a few studies have specifically examined tuberculosis mortality due to *M. bovis*. In at least two reports, it appeared to be higher in patients infected with this organism than *M. tuberculosis*; however, patient characteristics, including underlying illnesses, may not have been comparable. Two studies found case fatality rates of 10-14% in people with uncontrolled HIV infections and tuberculosis caused by *M. bovis*.

Groups at elevated risk of infection with *M. bovis* include farmers, abattoir workers, veterinary personnel, zoo staff and others who may handle infected animals or tissues, and those who drink unpasteurized milk. The incidence of tuberculosis due to this organism is poorly understood, as the causative organism is not always identified. Approximately 25-30% of all tuberculosis cases were thought to be caused by *M. bovis* before the advent of milk pasteurization and bovine tuberculosis control programs, at a time when close contact with livestock was also more common. Newer studies estimate that this organism is responsible for < 1% to 5-10%

or more of the tuberculosis cases in various countries, and usually < 2% of the cases in developed nations with good bovine tuberculosis programs (e.g., approximately 230 cases in the U.S. each year). In endemic areas, clinical cases are especially common in children, who are more susceptible than adults and may be exposed to *M. bovis* in unpasteurized milk products. However, most cases in developed countries now occur in elderly or immunosuppressed individuals, as the result of reactivated infections acquired in childhood, and in immigrants from endemic regions. Small clusters of *M. bovis* infections seem to be caused by person-to-person transmission, especially among people in close contact (e.g., families). Healthy people, as well as those who are immunocompromised, have been affected in these incidents.

Organisms other than M. bovis

M. caprae is uncommon in humans but it may represent a significant fraction of the zoonotic cases where *M. bovis* has been controlled. In Europe, this organism was responsible for <1% to 70% of the tuberculosis cases in various surveys. Risk factors seem to be similar to those for *M. bovis*, and both goats and cattle have been associated with human infections. Clinical cases caused by *M. microti* and *M. pinnipedii* are thought to be uncommon in humans. However, infections with *M. microti* may be underdiagnosed, as this organism is common in wild rodents in some areas, but it is difficult to culture and clinical suspicion is low.

Internet Resources

Canadian Food Inspection Agency. Bovine Tuberculosis
<https://www.inspection.gc.ca/animals/terrestrial-animals/diseases/reportable/bovine-tuberculosis/eng/1330205978967/1330206128556>

Centers for Disease Control and Prevention, United States. Tuberculosis
<https://www.cdc.gov/tb/default.htm>

European Centre for Disease Prevention and Control. Tuberculosis
<https://www.ecdc.europa.eu/en/tuberculosis>

European Union Reference Laboratory. Bovine Tuberculosis
<https://www.visavet.es/bovinetuberculosis/bovine-tb/eradication.php>

Michigan Bovine Tuberculosis Eradication Project
https://www.michigan.gov/emergingdiseases/0,4579,7-186-76711_78153-74719--,00.html

The Merck Manual
<http://www.merckmanuals.com/professional>

The Merck Veterinary Manual
<http://www.merckvetmanual.com>

Public Health Agency of Canada. Pathogen Safety Data Sheets and Risk Assessment
<https://www.canada.ca/en/public-health/services/laboratory-biosafety-biosecurity/pathogen-safety-data-sheets-risk-assessment.html>

World Organization for Animal Health (OIE).
<http://www.oie.int>

OIE Manual of Diagnostic Tests and Vaccines for Terrestrial Animals
<http://www.oie.int/international-standard-setting/terrestrial-manual/access-online/>

OIE Terrestrial Animal Health Code
<http://www.oie.int/international-standard-setting/terrestrial-code/access-online/>

United States Department of Agriculture, Animal and Plant Health Inspection Service (USDA APHIS) National Tuberculosis Eradication Program
<https://www.aphis.usda.gov/aphis/ourfocus/animalhealth/animal-disease-information/cattle-disease-information/national-tuberculosis-eradication-program>

World Health Organization. Tuberculosis
<https://www.who.int/tb/en/>

Acknowledgements

This factsheet was written by Anna Rovid Spickler, DVM, PhD, Veterinary Specialist from the Center for Food Security and Public Health. The U.S. Department of Agriculture Animal and Plant Health Inspection Service (USDA APHIS) provided funding for this factsheet through a series of cooperative agreements related to the development of resources for initial accreditation training.

The following format can be used to cite this factsheet. Spickler, Anna Rovid. 2019. Zoonotic Tuberculosis. Retrieved from <http://www.cfsph.iastate.edu/DiseaseInfo/factsheets.php>.

References

- Aiello SE, Moses MA, editors. The Merck veterinary manual. 11th ed. Kenilworth, NJ: Merck and Co; 2016. Tuberculosis and other mycobacterial infections. p. 687-92.
- Akkerman O, van der Loo K, Nijmeijer D, van der Werf T, Mulder B, Kremer K, van Soelingen D, van der Zanden A. *Mycobacterium bovis* infection in a young Dutch adult: transmission from an elderly human source? *Med Microbiol Immunol*. 2012;201(3):397-400.
- Albernaz TT, Oliveira CM, Lima DH, da Silva e Silva N, Cardoso DP, Lopes CT, Brito Mde F, da Silva JB, Salvarani FM, Leite RC, Barbosa JD. Comparison of the tuberculin test, histopathological examination, and bacterial culture for the diagnosis of tuberculosis (*Mycobacterium bovis*) in buffaloes (*Bubalus bubalis*) in Brazil. *Trop Anim Health Prod*. 2015;47(6):1153-9.
- Allen AR, Minozzi G, Glass EJ, Skuce RA, McDowell SW, Woolliams JA, Bishop SC. Bovine tuberculosis: the genetic basis of host susceptibility. *Proc Biol Sci*. 2010;277(1695):2737-45.
- Amato B, Capucchio TM, Biasibetti E, Mangano E, Boniotti BM, Pacciarini LM, Migliore S, Vitale M, Fiasconaro M, Di Marco Lo Presti V. Pathology and genetic findings in a rare case of *Mycobacterium caprae* infection in a sow. *Vet Microbiol*. 2017;205:71-4.
- Amato B, Mignacca SA, Pacciarini ML, Vitale M, Antoci S, Cucinotta S, Puleio R, Biasibetti E, Fiasconaro M, Capucchio MT, Di Marco Lo Presti V. An outbreak of bovine tuberculosis in a fallow deer herd (*Dama dama*) in Sicily. *Res Vet Sci*. 2016;106:116-20.
- Aranaz A, Liébana E, Gómez-Mampaso E, Galán JC, Cousins D, Ortega A, Blázquez J, Baquero F, Mateos A, Suárez G, Domínguez L. *Mycobacterium tuberculosis* subsp. *caprae* subsp. nov.: a taxonomic study of a new member of the *Mycobacterium tuberculosis* complex isolated from goats in Spain. *Int J Syst Bacteriol*. 1999;49 Pt 3:1263-73.
- Ameni G, Tafess K, Zewde A, Egualé T, Tilahun M, Hailu T, Sirak A, Salguero FJ, Berg S, Aseffa A, Hewinson RG, Vordermeier HM. Vaccination of calves with *Mycobacterium bovis* Bacillus Calmette-Guerin reduces the frequency and severity of lesions of bovine tuberculosis under a natural transmission setting in Ethiopia. *Transbound Emerg Dis*. 2018 ;65(1):96-104.
- Animal Health Australia. The National Animal Health Information System [NAHIS]. Bovine tuberculosis [online]. NAHIS; 2001 Oct. Available at: <http://www.aahc.com.au/nahis/disease/dislist.asp>. * Accessed 2 Nov 2001.
- Banos G, Winters M, Mrode R, Mitchell AP, Bishop SC, Woolliams JA, Coffey MP. Genetic evaluation for bovine tuberculosis resistance in dairy cattle. *J Dairy Sci*. 2017;100(2):1272-81.
- Barbier E, Rochelet M, Gal L, Boschirolu ML, Hartmann A. Impact of temperature and soil type on *Mycobacterium bovis* survival in the environment. *PLoS One*. 2017;12(4):e0176315.
- Barnett JE, Booth P, Brewer JI, Chanter J, Cooper T, Crawshaw T, Davison NJ, Greenwood A, Riley P, Smith NH, Wessels M. *Mycobacterium bovis* infection in a grey seal pup (*Halichoerus grypus*). *Vet Rec*. 2013;173(7):168.
- Barron MC, Pech RP, Whitford J, Yockney JJ, de Lisle GW, Nugent G. Longevity of *Mycobacterium bovis* in brushtail possum (*Trichosurus vulpecula*) carcasses, and contact rates between possums and carcasses. *N Z Vet J*. 2011;59(5): 209-17.
- Bennett AD, Lalor S, Schwarz T, Gunn-Moore DA. Radiographic findings in cats with mycobacterial infections. *J Feline Med Surg*. 2011;13(10):718-24.
- Best M, Sattar SA, Springthorpe VS, Kennedy ME. Efficacies of selected disinfectants against *Mycobacterium tuberculosis*. *J Clin Microbiol*. 1990;28(10):2234-9.
- Bezós J, Alvarez J, de Juan L, Romero B, Rodríguez S, Fernández-de-Mera IG, Hewinson RG, Vordermeier M, Mateos A, Domínguez L, Aranaz A. Assessment of *in vivo* and *in vitro* tuberculosis diagnostic tests in *Mycobacterium caprae* naturally infected caprine flocks. *Prev Vet Med*. 2011;100(3-4):187-92.

- Bezoz J, Alvarez J, Romero B, Aranaz A, Juan L. Tuberculosis in goats: assessment of current *in vivo* cell-mediated and antibody-based diagnostic assays. *Vet J.* 2012 ;191(2):161-5.
- Bezoz J, Casal C, Díez-Delgado I, Romero B, Liandris E, Álvarez J, Sevilla IA, Juan Ld, Domínguez L, Gortázar C. Goats challenged with different members of the *Mycobacterium tuberculosis* complex display different clinical pictures. *Vet Immunol Immunopathol.* 2015;167(3-4):185-9.
- Bezoz J, Casal C, Romero B, Schroeder B, Hardegger R, Raeber AJ, López L, Rueda P, Domínguez L. Current ante-mortem techniques for diagnosis of bovine tuberculosis. *Res Vet Sci.* 2014;97 Suppl:S44-52.
- Bezoz J, Marqués S, Álvarez J, Casal C, Romero B, Grau A, Mínguez O, Domínguez L, de Juan L. Evaluation of single and comparative intradermal tuberculin tests for tuberculosis eradication in caprine flocks in Castilla y León (Spain). *Res Vet Sci.* 2014;96(1):39-46.
- Bezoz J, Roya A, Infantes-Lorenzo JA, González I, Venteo A, Romero B, Grau A, Mínguez O, Domínguez Lucía de Juana L. The use of serological tests in combination with the intradermal tuberculin test maximizes the detection of tuberculosis infected goats. *Vet Immunol Immunopathol.* 2018;199:43-52.
- Biberstein EL, Holzworth J. Tuberculosis. In: Holzworth J, editor. *Diseases of the cat.* Philadelphia: WB Saunders; 1987. p. 284-6.
- Broughan JM, Crawshaw TR, Downs SH, Brewer J, Clifton-Hadley RS. *Mycobacterium bovis* infections in domesticated non-bovine mammalian species. Part 2: A review of diagnostic methods. *Vet J.* 2013;198(2):346-51.
- Broughan JM, Downs SH, Crawshaw TR, Upton PA, Brewer J, Clifton-Hadley RS. *Mycobacterium bovis* infections in domesticated non-bovine mammalian species. Part 1: Review of epidemiology and laboratory submissions in Great Britain 2004-2010. *Vet J.* 2013;198(2):339-45.
- Brüns AC, Tanner M, Williams MC, Botha L, O'Brien A, Fosgate GT, van Helden PD, Clarke J, Michel AL. Diagnosis and implications of *Mycobacterium bovis* infection in banded mongooses (*Mungos mungo*) in the Kruger National Park, South Africa. *J Wildl Dis.* 2017 ;53(1):19-29.
- Buddle BM, de Lisle GW, Griffin JF, Hutchings SA. Epidemiology, diagnostics, and management of tuberculosis in domestic cattle and deer in New Zealand in the face of a wildlife reservoir. *N Z Vet J.* 2015;63 Suppl 1:19-27.
- Buddle BM, Parlane NA, Wedlock DN, Heiser A. Overview of vaccination trials for control of tuberculosis in cattle, wildlife and humans. *Transbound Emerg Dis.* 2013;60 Suppl 1:136-46.
- Buddle BM, Vordermeier HM, Chambers MA(2), de Klerk-Lorist LM. Efficacy and safety of BCG vaccine for control of tuberculosis in domestic livestock and wildlife. *Front Vet Sci.* 2018;5:259.
- Busch F, Bannerman F, Liggett S, Griffin F, Clarke J, Lyashchenko KP, Rhodes S. Control of bovine tuberculosis in a farmed red deer herd in England. *Vet Rec.* 2017;180(3):68.
- Buss BF, Keyser-Metobo A, Rother J, Holtz L, Gall K, Jereb J, Murphy CN, Iwen PC, Robbe-Austerman S, Holcomb MA, Infield P. Possible airborne person-to-person transmission of *Mycobacterium bovis* - Nebraska 2014-2015. *MMWR Morb Mortal Wkly Rep.* 2016;65(8):197-201.
- Butler KL, Fitzgerald SD, Berry DE, Church SV, Reed WM, Kaneene JB. Experimental inoculation of European starlings (*Sturnus vulgaris*) and American crows (*Corvus brachyrhynchos*) with *Mycobacterium bovis*. *Avian Dis.* 2001;45:709-18.
- Byrom AE, Caley P, Paterson BM, Nugent G. Feral ferrets (*Mustela furo*) as hosts and sentinels of tuberculosis in New Zealand. *N Z Vet J.* 2015;63 Suppl 1:42-53.
- Campbell TA, Long DB, Bazan LR, Thomsen BV, Robbe-Austerman S, Davey RB, Soliz LA, Swafford SR, VerCauteren KC. Absence of *Mycobacterium bovis* in feral swine (*Sus scrofa*) from the southern Texas border region. *J Wildl Dis.* 2011;47(4):974-8.
- Cassidy JP. The pathogenesis and pathology of bovine tuberculosis with insights from studies of tuberculosis in humans and laboratory animal models. *Vet Microbiol.* 2006; 112 151-61.
- Cavanagh R, Begon M, Bennett M, Ergon T, Graham IM, De Haas PE, Hart CA, Koedam M, Kremer K, Lambin X, Roholl P, Soolingen Dv Dv. *Mycobacterium microti* infection (vole tuberculosis) in wild rodent populations. *J Clin Microbiol.* 2002;40(9):3281-5.
- Černá P, O'Halloran C, Sjatkovska J O, Gunn-Moore DA. Outbreak of tuberculosis caused by *Mycobacterium bovis* in a cattery of Abyssinian cats in Italy. *Transbound Emerg Dis.* 2019 ;66(1):250-8.
- Chambers MA. Review of the diagnosis and study of tuberculosis in non-bovine wildlife species using immunological methods. *Transbound Emerg Dis.* 2009;56(6-7):215-27.
- Christie LJ, Loeffler AM, Honarmand S, Flood JM, Baxter R, Jacobson S, Alexander R, Glaser CA. Diagnostic challenges of central nervous system tuberculosis. *Emerg Infect Dis.* 2008;14(9):1473-5.
- Churchyard G, Kim P, Shah NS, Rustomjee R, Gandhi N, Mathema B, Dowdy D, Kasmar A, Cardenas V. What we know about tuberculosis transmission: an overview. *J Infect Dis.* 2017;216(suppl_6):S629-S635.
- Clarke KA, Fitzgerald SD, Zwick LS, Church SV, Kaneene JB, Wismer AR, Bolin CA, Hattey JA, Yuzbasiyan-Gurkan V. Experimental inoculation of meadow voles (*Microtus pennsylvanicus*), house mice (*Mus musculus*), and Norway rats (*Rattus norvegicus*) with *Mycobacterium bovis*. *J Wildl Dis.* 2007;43:353-65.
- Clarke KR, Fitzgerald SD, Hattey JA, Bolin CA, Berry DE, Church SV, Reed WM. Experimental inoculation of wild turkeys (*Meleagris gallopavo*) with *Mycobacterium bovis*. *Avian Dis.* 2006;50:131-4.
- Corcoran JP, Hallifax RJ, Bettinson HV, Psallidas I, Rahman NM. Tuberculous pleuritis secondary to *Mycobacterium bovis* in a veterinarian. *Clin Respir J.* 2016;10(4):500-3.
- Corner LA. The role of wild animal populations in the epidemiology of tuberculosis in domestic animals: How to assess the risk. *Vet Microbiol.* 2006; 112:303-12.
- Corner LA, Gormley E, Pfeiffer DU. Primary isolation of *Mycobacterium bovis* from bovine tissues: conditions for maximising the number of positive cultures. *Vet Microbiol.* 2012;156(1-2):162-71.

- Corner LA, Murphy D, Gormley E. *Mycobacterium bovis* infection in the Eurasian badger (*Meles meles*): the disease, pathogenesis, epidemiology and control. *J Comp Pathol*. 2011;144(1):1-24.
- Cousins DV. *Mycobacterium bovis* infection and control in domestic livestock. *Rev Sci Tech*. 2001;20:71-85.
- Cousins DV, Bastida R, Cataldi A, Quse V, Redrobe S, et. al. Tuberculosis in seals caused by a novel member of the *Mycobacterium tuberculosis* complex: *Mycobacterium pinnipedii* sp. nov. *Int J Syst Evol Microbiol*. 2003;53(Pt 5):1305-14.
- Cousins DV, Florisson N. A review of tests available for use in the diagnosis of tuberculosis in non-bovine species. *Rev Sci Tech*. 2005;24:1039-59.
- Dalla Costa ER, Vasconcelos SE, Esteves LS, Gomes HM, Gomes LL, et al. Multidrug-resistant *Mycobacterium tuberculosis* of the Latin American Mediterranean lineage, wrongly identified as *Mycobacterium pinnipedii* (spoligotype international type 863 [SIT863]), causing active tuberculosis in South Brazil. *J Clin Microbiol*. 2015;53(12):3805-11.
- Dawson KL, Bell A, Kawakami RP, Coley K, Yates G, Collins DM. Transmission of *Mycobacterium orygis* (*M. tuberculosis* complex species) from a tuberculosis patient to a dairy cow in New Zealand. *J Clin Microbiol*. 2012;50(9):3136-8.
- Deforges L, Boulouis HJ, Thibaud JL, Boulouha L, Sougakoff W, Blot S, Hewinson G, Truffot-Pernot C, Haddad N. First isolation of *Mycobacterium microti* (Llama-type) from a dog. *Vet Microbiol*. 2004;103(3-4):249-53.
- De Garine-Wichatitsky M, Caron A, Kock R, Tschopp R, Munyeme M, Hofmeyr M, Michel A. A review of bovine tuberculosis at the wildlife-livestock-human interface in sub-Saharan Africa. *Epidemiol Infect*. 2013;141(7):1342-56.
- de Jong E, Rentenaar RJ, van Pelt R, de Lange W, Schreurs W, van Soelingen D, Sturm PD. Two cases of *Mycobacterium microti*-induced culture-negative tuberculosis. *J Clin Microbiol*. 2009;47(9):3038-40.
- de Kantor IN, LoBue PA, Thoen CO. Human tuberculosis caused by *Mycobacterium bovis* in the United States, Latin America and the Caribbean. *Int J Tuberc Lung Dis*. 2010;14(11):1369-73.
- de Kantor IN, Ritacco V. An update on bovine tuberculosis programmes in Latin American and Caribbean countries. *Vet Microbiol*. 2006;112:111-8.
- Delahay RJ, Smith GC, Barlow AM, Walker N, Harris A, Clifton-Hadley RS, Cheeseman CL. Bovine tuberculosis infection in wild mammals in the southwest region of England: a survey of prevalence and a semi-quantitative assessment of the relative risks to cattle. *Vet J*. 2007;173(2):287-301.
- de Val Pérez B, López-Soria S, Nofrarías M, Martín M, Vordermeier HM, Villarreal-Ramos B, Romera N, Escobar M, Solanes D, Cardona PJ, Domingo M. Experimental model of tuberculosis in the domestic goat after endobronchial infection with *Mycobacterium caprae*. *Clin Vaccine Immunol*. 2011;18(11):1872-81.
- Dias MF, Bernardes Filho F, Quaresma MV, Nascimento LV, Nery JA, Azulay DR. Update on cutaneous tuberculosis. *An Bras Dermatol*. 2014;89(6):925-38.
- Di Marco V, Mazzone P, Capucchio MT, Boniotti MB, Aronica V, Russo M, Fiasconaro M, Cifani N, Corneli S, Biasibetti E, Biagetti M, Pacciarini ML, Cagiola M, Pasquali P, Marianelli C. Epidemiological significance of the domestic black pig (*Sus scrofa*) in maintenance of bovine tuberculosis in Sicily. *J Clin Microbiol*. 2012;50(4):1209-18.
- Domingo M, Vidal E, Marco A. Pathology of bovine tuberculosis. *Res Vet Sci*. 2014;97 Suppl:S20-9.
- Drain PK, Bajema KL, Dowdy D, Dheda K, Naidoo K, Schumacher SG, Ma S, Meermeier E, Lewinsohn DM, Sherman DR. Incipient and subclinical tuberculosis: a clinical review of early stages and progression of infection. *Clin Microbiol Rev*. 2018;31(4). pii: e00021-18.
- Drewe JA, Foote AK, Sutcliffe RL, Pearce GP. Pathology of *Mycobacterium bovis* infection in wild meerkats (*Suricata suricatta*). *J Comp Pathol*. 2009 ;140(1):12-24.
- Dürr S, Müller B, Alonso S, Hattendorf J, Laisse CJ, van Helden PD, Zinsstag J. Differences in primary sites of infection between zoonotic and human tuberculosis: results from a worldwide systematic review. *PLoS Negl Trop Dis*. 2013;7(8):e2399.
- El-Sayed A, El-Shannat S, Kamel M, Castañeda-Vazquez MA, Castañeda-Vazquez H. Molecular epidemiology of *Mycobacterium bovis* in humans and cattle. *Zoonoses Public Health*. 2016;63(4):251-64.
- Eroksuz Y, Baydar E, Otlu B, Dabak M, Eroksuz H, Karabulut B, Incili CA, Timurkan MO. Case report: systemic tuberculosis caused by *Mycobacterium bovis* in a cat. *BMC Vet Res*. 2019;15(1):9.
- Etter E, Donado P, Jori F, Caron A, Goutard F, Roger F. Risk analysis and bovine tuberculosis, a re-emerging zoonosis. *Ann N Y Acad Sci*. 2006;1081:61-73.
- Evans JT, Smith EG, Banerjee A, Smith RM, Dale J, Innes JA, Hunt D, Tweddell A, Wood A, Anderson C, Hewinson RG, Smith NH, Hawkey PM, Sonnenberg P. Cluster of human tuberculosis caused by *Mycobacterium bovis*: evidence for person-to-person transmission in the UK. *Lancet*. 2007;369:1236-8.
- Faurholt-Jepsen D, Lillebaek T, Nielsen M-Y, Nielsen SD. *Mycobacterium bovis* meningitis in young Nigerian-born male. *Scand J Infect Dis*. 2014;46(10):732-4.
- Fico R, Mariacher A, Franco A, Eleni C, Ciarrocca E, Pacciarini ML, Battisti A. Systemic tuberculosis by *Mycobacterium bovis* in a free-ranging Marsica brown bear (*Ursus arctos marsicanus*): a case report. *BMC Vet Res*. 2019;15(1):152.
- Fine AE, Bolin CA, Gardiner JC, Kaneene JB. A study of the persistence of *Mycobacterium bovis* in the environment under natural weather conditions in Michigan, USA. *Vet Med Int*. 2011;2011:765430.
- Fine AE, O'Brien DJ, Winterstein SR, Kaneene JB. An effort to isolate *Mycobacterium bovis* from environmental substrates during investigations of bovine tuberculosis transmission sites (cattle farms and wildlife areas) in Michigan, USA. *ISRN Vet Sci*. 2011;2011:787181.
- Fink M, Schleicher C, Gonano M, Prodinger WM, Pacciarini M, Glawischnig W, Ryser-Degiorgis MP, Walzer C, Stalder GL, Lombardo D, Schobesberger H, Winter P, Büttner M. Red deer as maintenance host for bovine tuberculosis, Alpine region. *Emerg Infect Dis*. 2015;21(3):464-7.

- Fitzgerald SD, Boland KG, Clarke KR, Wismer A, Kaneene JB, Berry DE, Church SV, Hattey JA, Bolin CA. Resistance of Mallard ducks (*Anas platyrhynchos*) to experimental inoculation with *Mycobacterium bovis*. *Avian Dis*. 2005;49:144-6.
- Fitzgerald SD, Kaneene JB. Wildlife reservoirs of bovine tuberculosis worldwide: hosts, pathology, surveillance, and control. *Vet Pathol*. 2013;50(3):488-99.
- Fitzgerald SD, Schooley AM, Berry DE, Kaneene JB. Antimicrobial susceptibility testing of *Mycobacterium bovis* isolates from Michigan white-tailed deer during the 2009 hunting season. *Vet Med Int*. 2010;2011:903683.
- Fitzgerald SD, Zwick LS, Berry DE, Church SV, Kaneene JB, Reed WM. Experimental inoculation of pigeons (*Columba livia*) with *Mycobacterium bovis*. *Avian Dis*. 2003;47:470-5.
- Forbes BA, Hall GS, Miller MB, Novak SM, Rowlinson MC, Salfinger M, Somoskövi A, Warshauer DM, Wilson ML. Practice guidelines for clinical microbiology laboratories: *Mycobacteria*. *Clin Microbiol Rev*. 2018;31(2): e00038-17..
- Frankel A, Penrose C, Emer J. Cutaneous tuberculosis: a practical case report and review for the dermatologist. *J Clin Aesthet Dermatol*. 2009;2(10):19-27.
- Fritsche A, Engel R, Buhl D, Zellweger JP. *Mycobacterium bovis* tuberculosis: from animal to man and back. *Int J Tuberc Lung Dis*. 2004;8:903-4.
- García-Bocanegra I, Barranco I, Rodríguez-Gómez IM, Pérez B, Gómez-Laguna J, Rodríguez S, Ruiz-Villamayor E, Perea A. Tuberculosis in alpacas (*Lama pacos*) caused by *Mycobacterium bovis*. *J Clin Microbiol*. 2010;48(5):1960-4.
- García-Jiménez WL, Benítez-Medina JM, Fernández-Llario P, Abecia JA, García-Sánchez A, Martínez R, Risco D, Ortiz-Peláez A, Salguero FJ, Smith NH, Gómez L, Hermoso de Mendoza J. Comparative pathology of the natural infections by *Mycobacterium bovis* and by *Mycobacterium caprae* in wild boar (*Sus scrofa*). *Transbound Emerg Dis*. 2013;60(2):102-9.
- Garner G, Saville P, Fediaevsky A. Manual for the recognition of exotic diseases of livestock: A reference guide for animal health staff [online]. Food and Agriculture Organization of the United Nations [FAO]; 2004. Tuberculosis (bovine). Available at: <http://www.spc.int/rahs/>. * Accessed 9 Oct 2007.
- Gavier-Widén D, Cooke MM, Gallagher J, Chambers MA, Gortázar C. A review of infection of wildlife hosts with *Mycobacterium bovis* and the diagnostic difficulties of the 'no visible lesion' presentation. *N Z Vet J*. 2009;57(3):122-31.
- Ghodbane R, Mba Medie F, Lepidi H, Nappez C, Drancourt M. Long-term survival of tuberculosis complex mycobacteria in soil. *Microbiology*. 2014;160(Pt 3):496-501.
- González-Duarte A, Ponce de León A, Osornio JS. Importance of differentiating *Mycobacterium bovis* in tuberculous meningitis. *Neurol Int*. 2011;3(3):e9.
- Gormley E, Corner LA, Costello E, Rodriguez-Campos S. Bacteriological diagnosis and molecular strain typing of *Mycobacterium bovis* and *Mycobacterium caprae*. *Res Vet Sci*. 2014;97 Suppl:S30-43.
- Gormley E, Doyle M, Duignan A, Good M, More SJ, Clegg TA. Identification of risk factors associated with disclosure of false positive bovine tuberculosis reactors using the gamma-interferon (IFN γ) assay. *Vet Res*. 2013;44:117.
- Gortazar C, Vicente J, Boadella M, Ballesteros C, Galindo RC, Garrido J, Aranaz A, de la Fuente J. Progress in the control of bovine tuberculosis in Spanish wildlife. *Vet Microbiol*. 2011;151(1-2):170-8.
- Grange JM. *Mycobacterium bovis* infection in human beings. *Tuberculosis (Edinb)*. 2001;81:71-7.
- Greenwald R, Lyashchenko O, Esfandiari J, Miller M, Mikota S, Olsen JH, Ball R, Dumonceaux G, Schmitt D, Moller T, Payeur JB, Harris B, Sofranko D, Waters WR, Lyashchenko KP. Highly accurate antibody assays for early and rapid detection of tuberculosis in African and Asian elephants. *Clin Vaccine Immunol*. 2009;16(5):605-12.
- Gunn-Moore DA. Feline mycobacterial infections. *Vet J*. 2014;201(2):230-8.
- Gunn-Moore DA, McFarland SE, Brewer JI, Crawshaw TR, Clifton-Hadley RS, Kovalik M, Shaw DJ. Mycobacterial disease in cats in Great Britain: I. Culture results, geographical distribution and clinical presentation of 339 cases. *J Feline Med Surg*. 2011;13(12):934-44.
- Gunn-Moore DA, McFarland SE, Schock A, Brewer JI, Crawshaw TR, Clifton-Hadley RS, Shaw DJ. Mycobacterial disease in a population of 339 cats in Great Britain: II. Histopathology of 225 cases, and treatment and outcome of 184 cases. *J Feline Med Surg*. 2011;13(12):945-52.
- Hammer P, Richter E, Rüsck-Gerdes S, Walte HG, Matzen S, Kiesner C. Inactivation of *Mycobacterium bovis* ssp. *caprae* in high-temperature, short-term pasteurized pilot-plant milk. *J Dairy Sci*. 2015;98(3):1634-9.
- Hang'ombe MB, Munyeme M, Nakajima C, Fukushima Y, Suzuki H, Matandiko W, Ishii A, Mweene AS, Suzuki Y. *Mycobacterium bovis* infection at the interface between domestic and wild animals in Zambia. *BMC Vet Res*. 2012;8:221.
- Hines ME, Kreeger JM, Herron AJ. Mycobacterial infections of animals: pathology and pathogenesis. *Lab Anim Sci*. 1995;45(4):334-51.
- Hlavsa MC, Moonan PK, Cowan LS, Navin TR, Kammerer JS, Morlock GP, Crawford JT, Lobue PA. Human tuberculosis due to *Mycobacterium bovis* in the United States, 1995-2005. *Clin Infect Dis*. 2008;47(2):168-75.
- Hlokwe TM, Michel AL, Mitchel E, Gcebe N, Reininghaus B. First detection of *Mycobacterium bovis* infection in giraffe (*Giraffa camelopardalis*) in the Greater Kruger National Park complex: Role and implications. *Transbound Emerg Dis*. 2019 Jun 24 [Epub ahead of print].
- Hunter RL. Pathology of post primary tuberculosis of the lung: an illustrated critical review. *Tuberculosis (Edinb)*. 2011;91(6):497-509.
- Jenkins HE, Yuen CM, Rodriguez CA, Nathavitharana RR, McLaughlin MM, Donald P, Marais BJ, Becerra MC. Mortality in children diagnosed with tuberculosis: a systematic review and meta-analysis. *Lancet Infect Dis*. 2017;17(3): 285-95.
- Judge J, McDonald RA, Walker N, Delahay RJ. Effectiveness of biosecurity measures in preventing badger visits to farm buildings. *PLoS One*. 2011;6(12):e28941.
- Jurczynski K, Lyashchenko KP, Gomis D, Moser I, Greenwald R, Moisson P. Pinniped tuberculosis in Malayan tapirs (*Tapirus indicus*) and its transmission to other terrestrial mammals. *J Zoo Wildl Med*. 2011;42(2):222-7.

- Jurczynski K, Scharpegge J, Ley-Zaporozhan J, Ley S, Cracknell J, Lyashchenko K, Greenwald R, Schenk JP. Computed tomographic examination of South American sea lions (*Otaria flavescens*) with suspected *Mycobacterium pinnipedii* infection. *Vet Rec.* 2011;169(23):608-14.
- Kaneene JB, Miller R, Meyer RM. Abattoir surveillance: the U.S. experience. *Vet Microbiol.* 2006;112:273-82.
- Keating LA, Wheeler PR, Mansoor H, Inwald JK, Dale J, Hewinson RG, Gordon SV. The pyruvate requirement of some members of the *Mycobacterium tuberculosis* complex is due to an inactive pyruvate kinase: implications for *in vivo* growth. *Mol Microbiol.* 2005;56(1):163-74.
- Kelly P, Jahns H, Power E, Bainbridge J, Kenny K, Corpa JM, Cassidy JP, Callanan JJ. Mycobacteriosis in ostriches (*Struthio camelus*) due to infection with *Mycobacterium bovis* and *Mycobacterium avium* complex. *Avian Dis.* 2013;57(4):808-11.
- Kiers A, Klarenbeek A, Mendelts B, Van Soolingen D, Koëter G. Transmission of *Mycobacterium pinnipedii* to humans in a zoo with marine mammals. *Int J Tuberc Lung Dis.* 2008;12(12):1469-73.
- Kipar A, Burthe SJ, Hetzel U, Rokia MA, Telfer S, Lambin X, Birtles RJ, Begon M, Bennett M. *Mycobacterium microti* tuberculosis in its maintenance host, the field vole (*Microtus agrestis*): characterization of the disease and possible routes of transmission. *Vet Pathol.* 2014;51(5):903-14.
- Kovacs BJ, Chen YK, Kettering JD, Aprecio RM, Roy I. High-level disinfection of gastrointestinal endoscopes: are current guidelines adequate? *Am J Gastroenterol.* 1999;94(6):1546-50.
- Krajewska-Wędzina M, Augustynowicz-Kopec E, Weiner M, Szulowski K. Treatment for active tuberculosis in giraffe (*Giraffa camelopardalis*) in a zoo and potential consequences for public health - Case report. *Ann Agric Environ Med.* 2018;25(4):593-5.
- Krajewska-Wędzina M, Olech W, Kozińska M, Augustynowicz-Kopec E, Weiner M, Szulowski K. Bovine tuberculosis outbreak in farmed American bison (*Bison bison*) in Poland. *Pol J Vet Sci.* 2017;20(4):819-21.
- Kubica T, Rüsich-Gerdes S, Niemann S. *Mycobacterium bovis* subsp. *caprae* caused one-third of human *M. bovis*-associated tuberculosis cases reported in Germany between 1999 and 2001. *J Clin Microbiol.* 2003;41(7):3070-7.
- Lalor SM, Clarke S, Pink J, Parry A, Scurrill E, Fitzpatrick N, Watson F, O'Halloran C, Gunn-Moore D. Tuberculosis joint infections in four domestic cats. *JFMS Open Rep.* 2017;3(2):2055116917719401.
- Lan Z, Bastos M, Menzies D. Treatment of human disease due to *Mycobacterium bovis*: a systematic review. *Eur Respir J.* 2016;48(5):1500-3.
- Ledwon A, Szeleszczuk P, Zwolska Z, Augustynowicz-Kopec E, Sapierzynski R, Kozak M. Experimental infection of budgerigars (*Melopsittacus undulatus*) with five *Mycobacterium* species. *Avian Pathol.* 2008 ;37(1):59-64.
- Lee H, Kim JM, Jang Y, Lee K, Baek K, Lee B, Kim HY, Lee MH, Ryoo S, Bae YC, Choi EJ, So B. Bovine tuberculosis in an Asian small-clawed otter (*Aonyx cinerea*) in the Republic of Korea. *J Vet Diagn Invest.* 2015;27(5):651-5.
- Lees VW. Learning from outbreaks of bovine tuberculosis near Riding Mountain National Park: applications to a foreign animal disease outbreak. *Can Vet J.* 2004;45:28-34.
- le Roex N, Cooper D, van Helden PD, Hoal EG, Jolles AE. Disease control in wildlife: evaluating a test and cull programme for bovine tuberculosis in African buffalo. *Transbound Emerg Dis.* 2016;63(6):647-57.
- le Roex N, Koets AP, van Helden PD, Hoal EG. Gene polymorphisms in African buffalo associated with susceptibility to bovine tuberculosis infection. *PLoS One.* 2013;8(5):e64494.
- le Roex N, van Helden PD, Koets AP, Hoal EG. Bovine TB in livestock and wildlife: what's in the genes? *Physiol Genomics.* 2013;45(15):631-7.
- Li C, Liu L, Tao Y. Diagnosis and treatment of congenital tuberculosis: a systematic review of 92 cases. *Orphanet J Rare Dis.* 2019;14(1):131.
- Lin PL, Yee J, Klein E, Lerche NW. Immunological concepts in tuberculosis diagnostics for non-human primates: a review. *J Med Primatol.* 2008 ;37 Suppl 1:44-51.
- Lutze-Wallace C, Turcotte C, Glover G, Cousins D, Bell J, Berlie-Surujballi G, Barbeau Y, Randall G. Isolation of a *Mycobacterium microti*-like organism from a rock hyrax (*Procapra capensis*) in a Canadian zoo. *Can Vet J.* 2006;47(10):1011-3.
- Lyashchenko KP, Greenwald R, Esfandiari J, Meylan M, Burri IH, Zanolari P. Antibody responses in New World camelids with tuberculosis caused by *Mycobacterium microti*. *Vet Microbiol.* 2007;125(3-4):265-73.
- Maas M, Michel AL, Rutten VP. Facts and dilemmas in diagnosis of tuberculosis in wildlife. *Comp Immunol Microbiol Infect Dis.* 2013;36(3):269-85.
- Maas M, van Kooten PJ, Schreuder J, Morar D, Tijhaar E, Michel AL, Rutten VP. Development of a lion-specific interferon-gamma assay. *Vet Immunol Immunopathol.* 2012;149(3-4):292-7.
- Madeira S, Manteigas A, Ribeiro R, Otte J, Fonseca AP, Caetano P, Abernethy D, Boinas F. Factors that influence *Mycobacterium bovis* infection in red deer and wild boar in an epidemiological risk area for tuberculosis of game species in Portugal. *Transbound Emerg Dis.* 2017;64(3):793-804.
- Maggioli MF. A bloody evidence: Is *Mycobacterium bovis* bacteraemia frequent in cattle?! *Virulence.* 2016;7(7):748-50.
- Majoor CJ, Magis-Escurra C, van Ingen J, Boeree MJ, van Soolingen D. Epidemiology of *Mycobacterium bovis* disease in humans, The Netherlands, 1993-2007. *Emerg Infect Dis.* 2011;17(3):457-63.
- Major A, Holmes A, Warren-Smith C, Lalor S, Littler R, Schwarz T, Gunn-Moore D. Computed tomographic findings in cats with mycobacterial infection. *J Feline Med Surg.* 2016;18(6):510-7.
- Malama S, Johansen TB, Muma JB, Mwanza S, Djønne B, Godfroid J. Isolation and molecular characterization of *Mycobacterium bovis* from Kafue lechwe (*Kobus leche kafuensis*) from Zambia. *Trop Anim Health Prod.* 2014 ;46(1):153-7.
- Malone KM, Gordon SV. *Mycobacterium tuberculosis* complex members adapted to wild and domestic animals. *Adv Exp Med Biol.* 2017;1019:135-54.
- Mamo G, Bayleyegn G, Sisay Tessema T, Legesse M, Medhin G, Bjune G, Abebe F, Ameni G. Pathology of camel tuberculosis and molecular characterization of its causative agents in pastoral regions of Ethiopia. *PLoS One.* 2011;6(1):e15862.

- Maragou C, Theologie-Lygidakis N, Ioannidis P, Stenou A, Kanavaki S, Iatrou I, Tsofia MN. Primary tooth abscess caused by *Mycobacterium bovis* in an immunocompetent child. *Eur J Pediatr*. 2010;169(9):1143-5.
- Marcos LA, Spitzer ED, Mahapatra R, Ma Y, Halse TA, Shea J, Isabelle M, Lapierre P, Escuyer VE. *Mycobacterium orygis* lymphadenitis in New York, USA. *Emerg Infect Dis*. 2017;23(10):1749-51.
- Marianelli C, Cifani N, Capucchio MT, Fiasconaro M, Russo M, La Mancusa F, Pasquali P, Di Marco V. A case of generalized bovine tuberculosis in a sheep. *J Vet Diagn Invest*. 2010;22(3):445-8.
- Martín-Hernando MP, Torres MJ, Aznar J, Negro JJ, Gandía A, Gortázar C. Distribution of lesions in red and fallow deer naturally infected with *Mycobacterium bovis*. *J Comp Pathol*. 2010;142(1):43-50.
- Martins Melo A, Silva Filho RPD, von Groll A, Reis AJ, Diniz J, Perdigão J, Portugal I, da Silva PEA, Borelli Grecco F, Orzechowski Xavier M. Tuberculosis caused by *Mycobacterium pinnipedii* in a wild South American sea lion *Otaria flavescens* stranded in southern Brazil. *Dis Aquat Organ*. 2019;133(3):189-94.
- Mathews F, Macdonald DW, Taylor GM, Gelling M, Norman RA, Honess PE, Foster R, Gower CM, Varley S, Harris A, Palmer S, Hewinson G, Webster JP. Bovine tuberculosis (*Mycobacterium bovis*) in British farmland wildlife: the importance to agriculture. *Proc Biol Sci*. 2006;273(1584):357-65.
- Menin Á, Fleith R, Reck C, Marlow M, Fernandes P, Pilati C, Báfica A. Asymptomatic cattle naturally infected with *Mycobacterium bovis* present exacerbated tissue pathology and bacterial dissemination. *PLoS One*. 2013;8(1):e53884.
- Medeiros L, Marassi CD, Duarte RS, da Silva MG, Lilenbaum W. Comparison of decontamination methods for primary isolation of *Mycobacterium bovis* in paucibacillary bovine tissues. *Lett Appl Microbiol*. 2012;54(3):182-6.
- Menzies FD, Neill SD. Cattle-to-cattle transmission of bovine tuberculosis. *Vet J*. 2000;160:92-106.
- Michel AL, Müller B, van Helden PD. *Mycobacterium bovis* at the animal-human interface: a problem, or not? *Vet Microbiol*. 2010;140(3-4):371-81.
- Michelet L, de Cruz K, Phalente Y, Karoui C, Henault S, Beral M, Boschiroli ML. *Mycobacterium microti* infection in dairy goats, France. *Emerg Infect Dis*. 2016;22(3):569-70.
- Michelet L, de Cruz K, Zanella G, Aaziz R, Bulach T, Karoui C, Henault S, Joncour G, Boschiroli ML. Infection with *Mycobacterium microti* in animals in France. *J Clin Microbiol*. 2015;53(3):981-5.
- Miller M, Joubert J, Mathebula N, De Klerk-Lorist LM, Lyashchenko KP, Bengis R, van Helden P, Hofmeyr M, Olea-Popelka F, Greenwald R, Esfandiari J, Buss P. Detection of antibodies to tuberculosis antigens in free-ranging lions (*Panthera leo*) infected with *Mycobacterium bovis* in Kruger National Park, South Africa. *J Zoo Wildl Med*. 2012;43(2):317-23.
- Miller M, Michel A, van Helden P, Buss P. Tuberculosis in rhinoceros: an underrecognized threat? *Transbound Emerg Dis*. 2017;64(4):1071-8.
- Miller RS, Sweeney SJ. *Mycobacterium bovis* (bovine tuberculosis) infection in North American wildlife: current status and opportunities for mitigation of risks of further infection in wildlife populations. *Epidemiol Infect*. 2013;141(7):1357-70.
- Monies B, Jahans K, de la Rua R. Bovine tuberculosis in cats. *Vet Rec*. 2006;158:245-6.
- More SJ, Radunz B, Glanville RJ. Lessons learned during the successful eradication of bovine tuberculosis from Australia. *Vet Rec*. 2015;177(9):224-32.
- Muñoz Mendoza M, Juan Ld, Menéndez S, Ocampo A, Mourelo J, Sáez JL, Domínguez L, Gortázar C, García Marín JF, Balseiro A. Tuberculosis due to *Mycobacterium bovis* and *Mycobacterium caprae* in sheep. *Vet J*. 2012 ;191(2):267-9.
- Muñoz-Mendoza M, Marreros N, Boadella M, Gortázar C, Menéndez S, de Juan L, Bezos J, Romero B, Copano MF, Amado J, Sáez JL, Mourelo J, Balseiro A. Wild boar tuberculosis in Iberian Atlantic Spain: a different picture from Mediterranean habitats. *BMC Vet Res*. 2013;9:176.
- Muñoz-Mendoza M, Romero B, Del Cerro A, Gortazar C, Garcia-Marin JF, Menendez S, Mourelo J, de Juan L, Saez JL, Delahay RJ, Balseiro A. Sheep as a potential source of bovine TB: epidemiology, pathology and evaluation of diagnostic techniques. *Transbound Emerg Dis*. 2016;63:635-46.
- Murray A, Dineen A, Kelly P, McGoey K, Madigan G, NiGhallchoir E, Gunn-Moore DA. Nosocomial spread of *Mycobacterium bovis* in domestic cats. *J Feline Med Surg*. 2015 ;17(2):173-80.
- Narnaware SD, Dahiya SS, Tuteja FC, Nagarajan G, Nath K, Patil NV. Pathology and diagnosis of *Mycobacterium bovis* in naturally infected dromedary camels (*Camelus dromedarius*) in India. *Trop Anim Health Prod*. 2015;47(8):1633-6.
- Nebreda T, Álvarez-Prida E, Blanco B, Remacha MA, Samper S, Jiménez MS. Peritoneal tuberculosis due to *Mycobacterium caprae*. *IDCases*. 2016;4:50-2.
- Nelson JT, Orloski KA, Lloyd AL, Camacho M, Schoenbaum MA, Robbe-Austerman S, Thomsen BV, Hall SM. Evaluation of serodiagnostic assays for *Mycobacterium bovis* infection in elk, white-tailed deer, and reindeer in the United States. *Vet Med Int*. 2012;2012:563293.
- Ní Bhuachalla D, Corner LA, More SJ, Gormley E. The role of badgers in the epidemiology of *Mycobacterium bovis* infection (tuberculosis) in cattle in the United Kingdom and the Republic of Ireland: current perspectives on control strategies. *Vet Med (Auckl)*. 2014;6:27-38.
- Niemann S, Richter E, DalÄ¼gge-Tamm H, Schlesinger H, Graupner D, Königstein B, Gurath G, Greinert U, Rusch-Gerdes S. Two cases of *Mycobacterium microti* derived tuberculosis in HIV-negative immunocompetent patients. *Emerg Infect Dis*. 2000;6(5):539-42.
- Nigsch A, Glawischnig W, Bagó Z, Greber N. *Mycobacterium caprae* infection of red deer in western Austria-optimized use of pathology data to infer infection dynamics. *Front Vet Sci*. 2019;5:350.
- Nishi JS, Shury T, Elkin BT. Wildlife reservoirs for bovine tuberculosis (*Mycobacterium bovis*) in Canada: strategies for management and research. *Vet Microbiol*. 2006;112:325-38.
- Nugent G. Maintenance, spillover and spillback transmission of bovine tuberculosis in multi-host wildlife complexes: a New Zealand case study. *Vet Microbiol*. 2011;151(1-2):34-42.
- Nugent G, Buddle BM, Knowles G. Epidemiology and control of *Mycobacterium bovis* infection in brushtail possums (*Trichosurus vulpecula*), the primary wildlife host of bovine tuberculosis in New Zealand. *N Z Vet J*. 2015;63 Suppl 1:28-41.

- Nugent G, Gortazar C, Knowles G. The epidemiology of *Mycobacterium bovis* in wild deer and feral pigs and their roles in the establishment and spread of bovine tuberculosis in New Zealand wildlife. *N Z Vet J.* 2015;63 Suppl 1:54-67.
- Nugent G, Yockney IJ, Whitford EJ. Intraspecific transmission of *Mycobacterium bovis* among penned feral pigs in New Zealand. *J Wildl Dis.* 2011;47(2):364-72.
- O'Brien DJ, Fierke JS, Cooley TM, Fitzgerald SD, Cosgrove MK, Schmitt SM. Performance of diagnostic tests for bovine tuberculosis in North American furbearers and implications for surveillance. *Transbound Emerg Dis.* 2013;60 Suppl 1:67-73.
- O'Brien DJ, Schmitt SM, Berry DE, Fitzgerald SD, Lyon TJ, Vanneste JR, Cooley TM, Hogle SA, Fierke JS. Estimating the true prevalence of *Mycobacterium bovis* in free-ranging elk in Michigan. *J Wildl Dis.* 2008;44(4):802-10.
- O'Brien DJ, Schmitt SM, Fitzgerald SD, Berry DE. Management of bovine tuberculosis in Michigan wildlife: current status and near term prospects. *Vet Microbiol.* 2011;151(1-2):179-87.
- O'Brien DJ, Schmitt SM, Fitzgerald SD, Berry DE, Hickling GJ. Managing the wildlife reservoir of *Mycobacterium bovis*: the Michigan, USA, experience. *Vet Microbiol.* 2006;112:313-23.
- O'Brien DJ, Schmitt SM, Rudolph BA, Nugent G. Recent advances in the management of bovine tuberculosis in free-ranging wildlife. *Vet Microbiol.* 2011;151(1-2):23-33.
- O'Halloran C, Hope JC, Dobromylskij M, Burr P, McDonald K, Rhodes S, Roberts T, Dampney R, De la Rua-Domenech R, Robinson N, Gunn-Moore DA. An outbreak of tuberculosis due to *Mycobacterium bovis* infection in a pack of English Foxhounds (2016-2017). *Transbound Emerg Dis.* 2018;65(6):1872-84.
- O'Halloran C, Ioannidi O, Reed N, Murtagh K, Dettemering E, Van Poucke S, Gale J, Vickers J, Burr P, Gascoyne-Binzi D, Howe R, Dobromylskij M, Mitchell J, Hope J, Gunn-Moore D. Tuberculosis due to *Mycobacterium bovis* in pet cats associated with feeding a commercial raw food diet. *J Feline Med Surg.* 2019;21(8):667-81.
- Olea-Popelka F, Muwonge A, Perera A, Dean AS, Mumford E, Erlacher-Vindel E, Forcella S, Silk BJ, Ditiu L, El Idrissi A, Raviglione M, Cosivi O, LoBue P, Fujiwara PI. Zoonotic tuberculosis in human beings caused by *Mycobacterium bovis*- a call for action. *Lancet Infect Dis.* 2017;17(1):e21-e25.
- Orloski K, Robbe-Austerman S, Stuber T, Hench B, Schoenbaum M. Whole genome sequencing of *Mycobacterium bovis* isolated from livestock in the United States, 1989-2018. *Front Vet Sci.* 2018;5:253.
- Palmer MV. *Mycobacterium bovis*: characteristics of wildlife reservoir hosts. *Transbound Emerg Dis.* 2013;60 Suppl 1:1-13.
- Palmer MV, Waters WR. Advances in bovine tuberculosis diagnosis and pathogenesis: what policy makers need to know. *Vet Microbiol.* 2006;112:181-90.
- Panarella ML, Bimes RS. A naturally occurring outbreak of tuberculosis in a group of imported cynomolgus monkeys (*Macaca fascicularis*). *J Am Assoc Lab Anim Sci.* 2010;49(2):221-5.
- Panteix G, Gutierrez MC, Boschiroli ML, Rouviere M, Plaidy A, Pressac D, Porcheret H, Chyderiotis G, Ponsada M, Van Oortegem K, Salloum S, Cabuzel S, Bañuls AL, Van de Perre P, Godreuil S. Pulmonary tuberculosis due to *Mycobacterium microti*: a study of six recent cases in France. *J Med Microbiol.* 2010;59(Pt 8):984-9.
- Park D, Qin H, Jain S, Preziosi M, Minuto JJ, Mathews WC, Moser KS, Benson CA. Tuberculosis due to *Mycobacterium bovis* in patients coinfecting with human immunodeficiency virus. *Clin Infect Dis.* 2010;51(11):1343-6.
- Parte AC. LPSN - list of prokaryotic names with standing in nomenclature. (bacterio.net), 20 years on. *Int J Syst Evol Microbiol.* 68:1825-9; Genus *Mycobacterium*. Available at: <http://www.bacterio.net/mycobacterium.html>. Accessed 20 Sep 2019.
- Pate M, Svara T, Gombac M, Paller T, Zolnir-Dovc M, Emersic I, Prodinger WM, Bartos M, Zdovc I, Krt B, Pavlik I, Cvetnić Z, Pogacnik M, Ocepek M. Outbreak of tuberculosis caused by *Mycobacterium caprae* in a zoological garden. *J Vet Med B Infect Dis Vet Public Health.* 2006;53(8):387-92.
- Pate M, Zajc U, Kušar D, Žele D, Vengušt G, Pirš T, Ocepek M. *Mycobacterium* spp. in wild game in Slovenia. *Vet J.* 2016;208:93-5.
- Pavlik I, Trcka I, Parmova I, Svobodova J, Melicharek I, Nagy G, Cvetnic Z, Ocepek M, Pate M, Lipiec M. Detection of bovine and human tuberculosis in cattle and other animals in six Central European countries during the years 2000–2004. *Vet Med- Czech.* 2005;50 (7):291-9.
- Pedersen K, Miller RS, Anderson TD, Pabilonia KL, Lewis JR, Mihalco RL, Gortázar C, Gidlewski T. Limited antibody evidence of exposure to *Mycobacterium bovis* in feral swine (*Sus scrofa*) in the USA. *J Wildl Dis.* 2017 ;53(1):30-6.
- Pérez-Lago L, Navarro Y, García-de-Viedma D. Current knowledge and pending challenges in zoonosis caused by *Mycobacterium bovis*: a review. *Res Vet Sci.* 2014;97 Suppl:S94-S100.
- Pesciaroli M, Alvarez J, Boniotti MB, Cagiola M, Di Marco V, Marianelli C, Pacciarini M, Pasquali P. Tuberculosis in domestic animal species. *Res Vet Sci.* 2014;97 Suppl:S78-85.
- Pfeiffer DU. Epidemiology caught in the causal web of bovine tuberculosis. *Transbound Emerg Dis.* 2013;60 Suppl 1:104-10.
- Pollock C. Mycobacterial infection in the ferret. *Vet Clin Exot Anim.* 2012;15:121-9.
- Pollock JM, Rodgers JD, Welsh MD, McNair J. Pathogenesis of bovine tuberculosis: the role of experimental models of infection. *Vet Microbiol.* 2006;112:141-50.
- Prodinger WM, Eigentler A, Allerberger F, Schönbauer M, Glawischnig W. Infection of red deer, cattle, and humans with *Mycobacterium bovis* subsp. *caprae* in western Austria. *J Clin Microbiol.* 2002;40(6):2270-2.
- Prodinger WM, Indra A, Koksalan OK, Kilicaslan Z, Richter E. *Mycobacterium caprae* infection in humans. *Expert Rev Anti Infect Ther.* 2014;12(12):1501-13.

- Public Health Agency of Canada (PHAC). Pathogen Safety Data Sheets: infectious substances – *Mycobacterium tuberculosis* and *Mycobacterium tuberculosis* complex. Pathogen Regulation Directorate, PHAC; 2010 Nov. Available at: <https://www.canada.ca/en/public-health/services/laboratory-biosafety-biosecurity/pathogen-safety-data-sheets-risk-assessment/mycobacterium-tuberculosis-complex.html>. Accessed 20 Sept 2019.
- Rahim Z, Thapa J, Fukushima Y, van der Zanden AGM, Gordon SV, Suzuki Y, Nakajima C. Tuberculosis caused by *Mycobacterium orygis* in dairy cattle and captured monkeys in Bangladesh: a new scenario of tuberculosis in South Asia. *Transbound Emerg Dis*. 2017;64(6):1965-9.
- Ramdas KE, Lyashchenko KP, Greenwald R, Robbe-Austerman S, McManis C, Waters WR. *Mycobacterium bovis* infection in humans and cats in same household, Texas, USA, 2012. *Emerg Infect Dis*. 2015;21(3):480-3.
- Réveillaud É, Desvaux S, Boschirolu ML, Hars J, Faure É, Fediaevsky A, Cavalerie L, Chevalier F, Jabert P, Poliak S, Tourette I, Hendriks P, Richomme C. Infection of wildlife by *Mycobacterium bovis* in France assessment through a national surveillance system, Sylvatub. *Front Vet Sci*. 2018;5:262.
- Reviriego Gordejo FJ, Vermeersch JP. Towards eradication of bovine tuberculosis in the European Union. *Vet Microbiol*. 2006;112:101-9.
- Rhodes S, Holder T, Clifford D, Dexter I, Brewer J, Smith N, Waring L, Crawshaw T, Gillgan S, Lyashchenko K, Lawrence J, Clarke J, de la Rua-Domenech R, Vordermeier M. Evaluation of gamma interferon and antibody tuberculosis tests in alpacas. *Clin Vaccine Immunol*. 2012;19(10):1677-83.
- Riojas MA, McGough KJ, Rider-Riojas CJ, Rastogi N, Hazbón MH. Phylogenomic analysis of the species of the *Mycobacterium tuberculosis* complex demonstrates that *Mycobacterium africanum*, *Mycobacterium bovis*, *Mycobacterium caprae*, *Mycobacterium microti* and *Mycobacterium pinnipedii* are later heterotypic synonyms of *Mycobacterium tuberculosis*. *Int J Syst Evol Microbiol*. 2018;68(1):324-32.
- Rivière J, Carabin K, Le Strat Y, Hendriks P, Dufour B. Bovine tuberculosis surveillance in cattle and free-ranging wildlife in EU Member States in 2013: a survey-based review. *Vet Microbiol*. 2014;173(3-4):323-31.
- Roberts T, O'Connor C, Nuñez-García J, de la Rua-Domenech R, Smith NH. Unusual cluster of *Mycobacterium bovis* infection in cats. *Vet Rec*. 2014;174(13):326.
- Rodríguez E, Sánchez LP, Pérez S, Herrera L, Jiménez MS, Samper S, Iglesias MJ. Human tuberculosis due to *Mycobacterium bovis* and *M. caprae* in Spain, 2004-2007. *Int J Tuberc Lung Dis*. 2009;13(12):1536-41.
- Rodríguez S, Bezos J, Romero B, de Juan L, Álvarez J, Castellanos E, Moya N, Lozano F, Javed MT, Sáez-Llorente JL, Liébana E, Mateos A, Domínguez L, Aranaz A; Spanish Network on Surveillance and Monitoring of Animal Tuberculosis. *Mycobacterium caprae* infection in livestock and wildlife, Spain. *Emerg Infect Dis*. 2011;17(3):532-5.
- Rodríguez-Campos S, Smith NH, Boniotti MB, Aranaz A. Overview and phylogeny of *Mycobacterium tuberculosis* complex organisms: Implications for diagnostics and legislation of bovine tuberculosis. *Res Vet Sci*. 2014; 97:S5-S19.
- Roe WD, Lenting B, Kokosinska A, Hunter S, Duignan PJ, Gartrell B, Rogers L, Collins DM, de Lisle GW, Gedye K, Price-Carter M. Pathology and molecular epidemiology of *Mycobacterium pinnipedii* tuberculosis in native New Zealand marine mammals. *PLoS One*. 2019;14(2):e0212363.
- Roos EO, Olea-Popelka F, Buss P, Hausler GA, Warren R, van Helden PD, Parsons SDC, de Klerk-Lorist LM, Miller MA. Measuring antigen-specific responses in *Mycobacterium bovis*-infected warthogs (*Phacochoerus africanus*) using the intradermal tuberculin test. *BMC Vet Res*. 2018;14(1):360.
- Rüfenacht S, Bögli-Stuber K, Bodmer T, Jaunin VF, Jmaa DC, Gunn-Moore DA. *Mycobacterium microti* infection in the cat: a case report, literature review and recent clinical experience. *J Feline Med Surg*. 2011;13(3):195-204.
- Ryan TJ, Livingstone PG, Ramsey DS, de Lisle GW, Nugent G, Collins DM, Buddle BM. Advances in understanding disease epidemiology and implications for control and eradication of tuberculosis in livestock: the experience from New Zealand. *Vet Microbiol*. 2006;112:211-9.
- Salvador LCM, O'Brien DJ, Cosgrove MK, Stuber TP, Schooley AM, Crispell J, Church SV, Gröhn YT, Robbe-Austerman S, Kao RR. Disease management at the wildlife-livestock interface: Using whole-genome sequencing to study the role of elk in *Mycobacterium bovis* transmission in Michigan, USA. *Mol Ecol*. 2019;28(9):2192-205.
- Sánchez FD, Yela IJ, Alfonseca E, Campuzano J, Morales E, Aguilar C. Respiratory tract infection caused by *Mycobacterium bovis* in a black swan (*Cygnus atratus*). *Avian Pathol*. 2016;45(1):126-31.
- Sanchez J, Tomás L, Ortega N, Buendía AJ, del Rio L, Salinas J, Bezos J, Caro MR, Navarro JA. Microscopical and immunological features of tuberculoid granulomata and cavitary pulmonary tuberculosis in naturally infected goats. *J Comp Pathol*. 2011;145(2-3):107-17.
- Sarradell JE, Alvarez J, Biscia M, Zumarraga M, Wunschmann A, Armien AG, Perez AM. *Mycobacterium bovis* infection in a horse with granulomatous enterocolitis. *J Vet Diagn Invest*. 2015;27(2):203-5.
- Schaftenaar W, Lécú A, Greenwald R, Lyashchenko KP. Retrospective serological investigation of bovine tuberculosis in two gemsbok (*Oryx gazelle gazelle*) and an onager (*Equus hemionus onager*). *J Zoo Wildl Med*. 2013;44(4):1036-42.
- Schiller I, Waters WR, Vordermeier HM, Jemmi T, Welsh M, et al. Bovine tuberculosis in Europe from the perspective of an officially tuberculosis free country: trade, surveillance and diagnosis. *Vet Microbiol*. 2011;151(1-2):153-9.
- Schoepf K, Proding WM, Glawischnig W, Hofer E, Revilla-Fernandez S, Hofrichter J, Fritz J, Köfer J, Schmoll F. A two-years' survey on the prevalence of tuberculosis caused by *Mycobacterium caprae* in red deer (*Cervus elaphus*) in the Tyrol, Austria. *ISRN Vet Sci*. 2012;2012:245138.
- Schöning JM, Cerny N, Prohaska S, Wittenbrink MM, Smith NH, Bloemberg G, Pewsner M, Schiller I, Origi FC, Rysler-Degiorgis MP. Surveillance of bovine tuberculosis and risk estimation of a future reservoir formation in wildlife in Switzerland and Liechtenstein. *PLoS One*. 2013;8(1):e54253.
- Scott C, Cavanaugh JS, Silk BJ, Ershova J, Mazurek GH, LoBue PA, Moonan PK. Comparison of sputum-culture conversion for *Mycobacterium bovis* and *M. tuberculosis*. *Emerg Infect Dis*. 2017;23(3):456-62.

- Shivaprasad HL, Palmieri C. Pathology of mycobacteria in birds. *Vet Clin North Am Exot Anim Pract.* 2012;15:41-55.
- Shrikrishna D, de la Rua-Domenech R, Smith NH, Colloff A, Coutts I. Human and canine pulmonary *Mycobacterium bovis* infection in the same household: re-emergence of an old zoonotic threat? *Thorax.* 2009 ;64(1):89-91.
- Shury TK, Bergeson D. Lesion distribution and epidemiology of *Mycobacterium bovis* in elk and white-tailed deer in southwestern Manitoba, Canada. *Vet Med Int.* 2011;2011:591980.
- Shury TK, Bergeson D, Surujballi O, Lyashchenko KP, Greenwald R. Field evaluation of three blood-based assays for elk (*Cervus canadensis*) naturally infected with *Mycobacterium bovis*. *Prev Vet Med.* 2014;115(3-4):109-21.
- Stavinohova R, O'Halloran C, Newton JR, Oliver JAC, Scurrell E, Gunn-Moore DA. Feline ocular mycobacteriosis: clinical presentation, histopathological features, and outcome. *Vet Pathol.* 2019;56(5):749-60.
- Sunder S, Lanotte P, Godreuil S, Martin C, Boschirola ML, Besnier JM. Human-to-human transmission of tuberculosis caused by *Mycobacterium bovis* in immunocompetent patients. *J Clin Microbiol.* 2009;47(4):1249-51.
- Sweetline Anne N, Ronald BS, Kumar TM, Kannan P, Thangavelu A. Molecular identification of *Mycobacterium tuberculosis* in cattle. *Vet Microbiol.* 2017 ;198:81-7.
- Sweetline Anne N, Ronald BSM, Senthil Kumar TMA, Thangavelu A. Conventional and molecular determination of drug resistance in *Mycobacterium tuberculosis* and *Mycobacterium bovis* isolates in cattle. *Tuberculosis.* 2019;114:113-8.
- Swift BM, Convery TW, Rees CE. Evidence of *Mycobacterium tuberculosis* complex bacteraemia in intradermal skin test positive cattle detected using phage-RPA. *Virulence* 2016; 7(7):779-88.
- Thapa J, Nakajima C, Maharjan B, Poudell A, Suzuki Y. Molecular characterization of *Mycobacterium orygis* isolates from wild animals of Nepal. *Jpn J Vet Res.* 2015;63(3):151-8.
- Thapa J, Paudel S, Sadaula A, Shah Y, Maharjan B, Kaufman GE, McCauley D, Gairhe KP, Tsubota T, Suzuki Y, Nakajima C. *Mycobacterium orygis*-associated tuberculosis in free-ranging rhinoceros, Nepal, 2015. *Emerg Infect Dis.* 2016;22(3):570-2.
- Thorel MF, Karoui C, Varnerot A, Fleury C, Vincent V. Isolation of *Mycobacterium bovis* from baboons, leopards and a sea-lion. *Vet Res.* 1998;29(2):207-12.
- Torres-Gonzalez P, Soberanis-Ramos O, Martinez-Gamboa A, Chavez-Mazari B, Barrios-Herrera MT, Torres-Rojas M, Cruz-Hervert LP, Garcia-Garcia L, Singh M, Gonzalez-Aguirre A, Ponce de Leon-Garduño A, Sifuentes-Osornio J, Bobadilla-Del-Valle M. Prevalence of latent and active tuberculosis among dairy farm workers exposed to cattle infected by *Mycobacterium bovis*. *PLoS Negl Trop Dis.* 2013;7(4):e2177.
- Twomey DF, Crawshaw TR, Foster AP, Higgins RJ, Smith NH, Wilson L, McDean K, Adams JL, de la Rua-Domenech R. Suspected transmission of *Mycobacterium bovis* between alpacas. *Vet Rec.* 2009;165(4):121-2.
- U.K. Department for Environment Food and Rural Affairs [DEFRA] The Independent Scientific Group on Cattle TB (ISG). Pathogenesis and diagnosis of infections with *M. bovis* in cattle (Appendix C) [online]. DEFRA; 2003 Aug. Available at: <http://www.defra.gov.uk/animalh/tb/isg/report/annexc.htm>. * Accessed 11 Oct 2007.
- U.S. Department of Agriculture, Animal and Plant Health Inspection Service [USDA APHIS]. Bovine tuberculosis [online]. USDA APHIS; 1995 Sept. Available at: <http://www.aphis.usda.gov/oa/pubs/fsbtb.html>. * Accessed 5 Nov 2001.
- van der Burgt G. *Mycobacterium bovis* causing clinical disease in adult sheep. *Vet Rec.* 2010;166(10):306.
- van der Burgt GM, Crawshaw T, Foster AP, Denny DJ, Schock A. *Mycobacterium bovis* infection in dogs. *Vet Rec.* 2009;165(21):634.
- van der Merwe M, Bekker JL, van der Merwe P, Michel AL. Cooking and drying as effective mechanisms in limiting the zoonotic effect of *Mycobacterium bovis* in beef. *J S Afr Vet Assoc.* 2009;80(3):142-5.
- van Ingen J, Boeree MJ, Wright A, van der Laan T, Dekhuijzen PN, van Soolingen D. Second-line drug resistance in multidrug-resistant tuberculosis cases of various origins in the Netherlands. *Int J Tuberc Lung Dis.* 2008;12(11):1295-9.
- van Ingen J, Brosch R, van Soolingen D. Characterization of *Mycobacterium orygis*. *Emerg Infect Dis.* 2013;19(3):521-2.
- van Soolingen D, van der Zanden AGM, de Haas PEW, Noordhoek GT, Kiers A, Foudraïne NA, Portaels F, Kolk AHJ, Kremer K, van Embden JDA. Diagnosis of *Mycobacterium microti* infections among humans by using novel genetic markers. *J Clin Microbiol.* 1998;36:1840-5.
- van Zyl-Smit RN, Binder A, Meldau R, Mishra H, Semple PL, Theron G, Peter J, Whitelaw A, Sharma SK, Warren R, Bateman ED, Dheda K. Comparison of quantitative techniques including Xpert MTB/RIF to evaluate mycobacterial burden. *PLoS One.* 2011;6(12):e28815.
- Vayr F, Martin-Blondel G, Savall F, Soulat JM, Deffontaines G, Herin F. Occupational exposure to human *Mycobacterium bovis* infection: A systematic review. *PLoS Negl Trop Dis.* 2018;12(1):e0006208.
- Vazquez-Chacon CA, Martínez-Guarneros A, Couvin D, González-Y-Merchand JA, Rivera-Gutierrez S, Escobar-Gutierrez A, De-la-Cruz López JJ, Gomez-Bustamante A, Gonzalez-Macal GA, Gonçalves Rossi LM, Muñoz-Salazar R, Rastogi N, Vaughan G. Human multidrug-resistant *Mycobacterium bovis* infection in Mexico. *Tuberculosis (Edinb).* 2015;95(6):802-9.
- Viljoen IM, Sylvester TT, Parsons SDC, Millar RP, Helden PDV, Miller MA. Performance of the tuberculin skin test in *Mycobacterium bovis*-exposed and -unexposed African lions (*Panthera leo*). *J Wildl Dis.* 2019;55(3):537-43.
- Viljoen IM, van Helden PD, Millar RP. *Mycobacterium bovis* infection in the lion (*Panthera leo*): Current knowledge, conundrums and research challenges. *Vet Microbiol.* 2015;177(3-4):252-60.
- Vordermeier HM, Jones GJ, Buddle BM, Hewinson RG, Villarreal-Ramos B. Bovine tuberculosis in cattle: vaccines, DIVA tests, and host biomarker discovery. *Annu Rev Anim Biosci.* 2016;4:87-109.

Bovine Tuberculosis

- Vordermeier HM, Pérez de Val B, Buddle BM, Villarreal-Ramos B, Jones GJ, Hewinson RG, Domingo M. Vaccination of domestic animals against tuberculosis: review of progress and contributions to the field of the TBSTEP project. *Res Vet Sci.* 2014;97 Suppl:S53-60.
- Walter WD, Anderson CW, Smith R, Vanderklok M, Averill JJ, Vercauteren KC. On-farm mitigation of transmission of tuberculosis from white-tailed deer to cattle: literature review and recommendations. *Vet Med Int.* 2012;2012:616318.
- Warburton B, Livingstone P. Managing and eradicating wildlife tuberculosis in New Zealand. *N Z Vet J.* 2015;63 Suppl 1:77-88.
- Waters WR, Palmer MV. *Mycobacterium bovis* infection of cattle and white-tailed deer: Translational research of relevance to human tuberculosis. *ILAR J.* 2015;56(1):26-43.
- Wilkins MJ, Meyerson J, Bartlett PC, Spieldenner SL, Berry DE, Mosher LB, Kaneene JB, Robinson-Dunn B, Stobierski MG, Boulton ML. Human *Mycobacterium bovis* infection and bovine tuberculosis outbreak, Michigan, 1994-2007. *Emerg Infect Dis.* 2008;14(4):657-60.
- Witmer G, Fine AE, Gionfriddo J, Pipas M, Shively K, Piccolo K, Burke P. Epizootiologic survey of *Mycobacterium bovis* in wildlife and farm environments in northern Michigan. *J Wildl Dis.* 2010;46(2):368-78.
- Wobeser G. Bovine tuberculosis in Canadian wildlife: an updated history. *Can Vet J.* 2009;50(11):1169-76.
- Woodroffe R, Donnelly CA, Ham C, Jackson SY, Moyes K, Chapman K, Stratton NG, Cartwright SJ. Badgers prefer cattle pasture but avoid cattle: implications for bovine tuberculosis control. *Ecol Lett.* 2016;19(10):1201-8.
- World Organization for Animal Health (OIE). Press release: Update on wildlife diseases. OIE: 2000 Jan. Available at: http://www.oie.int/eng/press/A_000104.htm. * Accessed 9 Oct 2007.
- World Organization for Animal Health (OIE). Handistatus II [database online]. OIE; 2004. Available at: <http://www.oie.int/hs2/report.asp?lang=en> . Accessed 11 Oct 2007.
- World Organization for Animal Health (OIE) . Manual of diagnostic tests and vaccines for terrestrial animals [online]. Paris: OIE; 2018. Bovine tuberculosis (adopted 2009). Available at: https://www.oie.int/fileadmin/Home/eng/Health_standards/tahm/3.04.06_BOVINE_TB.pdf. Accessed 17 Sept 2019.
- World Organization for Animal Health (OIE). World animal health information database (WAHIS) [database online]. Bovine tuberculosis: Paris:OIE;2019. Available at: https://www.oie.int/wahis_2/public/wahid.php/Diseaseinformation/diseasehome. Accessed 17 Sept 2019.
- Xavier Emmanuel F, Seagar AL, Doig C, Rayner A, Claxton P, Laurenson I. Human and animal infections with *Mycobacterium microti*, Scotland. *Emerg Infect Dis.* 2007;13(12):1924-7.
- Yoshida S, Suga S, Ishikawa S, Mukai Y, Tsuyuguchi K, Inoue Y, Yamamoto T, Wada T. *Mycobacterium caprae* infection in captive Borneo elephant, Japan. *Emerg Infect Dis.* 2018;24(10):1937-40.
- Zanella G, Durand B, Hars J, Moutou F, Garin-Bastuji B, Duvauchelle A, Fermé M, Karoui C, Boschirolu ML. *Mycobacterium bovis* in wildlife in France. *J Wildl Dis.* 2008;44(1):99-108.
- Zanolari P, Robert N, Lyashchenko KP, Pfyffer GE, Greenwald R, Esfandiari J, Meylan M. Tuberculosis caused by *Mycobacterium microti* in South American camelids. *J Vet Intern Med.* 2009;23(6):1266-72.

*Link is defunct